

NATIONAL EMERGENCY OBSTETRIC AND NEONATAL CARE NEEDS ASSESSMENT

PROVINCIAL REPORT WESTERN PROVINCE 2012



Family Health Bureau



World Health
Organization



NATIONAL EMERGENCY OBSTETRIC AND NEONATAL CARE NEEDS ASSESSMENT

PROVINCIAL REPORT

WESTERN PROVINCE 2012



Family Health Bureau



National Emergency Obstetric and Neonatal Care Needs Assessment,
Provincial Report - Western Province 2012

Published in January 2014 by

Family Health Bureau,

Ministry of Health,

231, De Saram Place,

Colombo 10,

Sri Lanka

Web site: www.familyhealth.gov.lk

Printed by Lazergraphic (PVT) Ltd

ISBN: 978-955-1503-17-8

All rights reserved- reproduction of any part of the text is authorised, except for commercial purpose, provided the relevant authours and the Family Health Bureau, Ministry of Health, Sri Lanka, are acknowledged. No photographs may be reproduced, stored in a retrieval system or transmitted in any form without prior permission from the copyright holder.

This report is dedicated to

**The mothers in Sri Lanka
and
those who care for them;
past present and future**

Emergency Obstetric and Neonatal Care Needs Assessment Research team

Dr. Lalini Rajapaksa	Public Health Specialist
Dr. Lakshman Senanayake	Consultant Obstetrician and Gynecologists
Dr. Chandani Galwaduge	National Programme officer UNFPA
Dr. Dhammica Rowel	Consultant Community Physician
Dr. Nilmini Hemachandra	Consultant Community Physician
Dr. Padmal de Silva	Senior Registrar, Community Medicine

CORE GROUP

1. Dr. Terrance de Silva	Deputy Director General/ Medical Services
2. Dr. L. Siyabalagoda	Deputy Director General / Public Health Services
3. Dr. S. A. K. Gamage	Director/ Medical Services
4. Dr. S. Sathumugan	Director/ Tertiary /Care Services
5. Dr. Deepthi Perera	Director/Maternal and Child Health
6. Prof. Lalini Rajapaksha	Public Health Specialist
7. Dr. Lakshman Senanayake	Consultant Obstetric & Gynaecologist
8. Dr. Medha Weerasekera	Consultant Paediatrician
9. Dr. A. L. A. L Padmasri	RDHS/ Gampaha
10. Dr. Chithramalee de Silva	Deputy Director/ Maternal and child Health
11. Dr. C.I. Gunasekara	Deputy Director/ Castle Street Hospital for Women
12. Dr. Nilmini Hemachandra	Consultant Community Physician, FHB
13. Dr. Dhammica Rowel	Consultant Community Physician, FHB
14. Dr. Kapila Jayarathne	Consultant Community Physician, FHB
15. Dr. Nirosha Lansakara	Consultant Community Physician, FHB
16. Dr. Padmal de Silva	Senior Registrar/ Community Medicine
17. Dr. Chandani Galwaduge	National Programme Officer, UNFPA
18. Dr. Anoma Jayathilake	National Professional Officer, WHO
19. Dr. Nishamani Karawita	National Health Officer, UNICEF
20. Dr. Ananda Ranathunga	President, SLCOG (2011)
21. Dr. Hemantha Perera	President, SLCOG (2012)
22. Dr. Deepthi Samarage	President, Sri Lanka College of Pediatricians (2011)
23. Dr. Piyasena Samarakoon	Senior Assistant Secretary, Ministry of Health
24. Dr. Lakshmi Somathunga	DDG/ MS I
25. Dr. R. Balasooriya	D/TCS
26. Dr. L. Panapitiya	D/MS
27. Dr. P. Atapattu	D/TCS
28. Dr. K. Wickramasinghe	D/CSHW
29. Prof. Hemantha Senanayake	President, SLCOG (2013)
30. Prof. Asvini Fernando	President, Sri Lanka College of Pediatricians(2012)
31. Dr. Sri Lal de Silva	President, Sri Lanka College of Pediatricians (2013)
32. Dr. Deepika Attygalle	Health and Nutrition Specialist, UNICEF (since 2013)

Message from the Minister of Health



As a country, Sri Lanka is moving from lower middle income country to middle income country under the guidance of the Development Policy Framework of the Government of Sri Lanka, Mahinda Chinthana Vision for the Future. The objective of the government is to provide highest attainable standard of health for every human being that includes provision of access to timely acceptable health care of appropriate quality. Sri Lanka is always considered a role model as a middle income country that has successfully reduced maternal, infant and child mortality to levels comparable with those of developed countries. I am informed that further improvements in health status need more stringent attention in provision of quality health care. Also it is important to focus on patient's rights, patient satisfaction, while providing high quality of health services.

Each year, the government of Sri Lanka spends billions of rupees for the development of health services in the country. The benefits of this should reach every corner of the country. Therefore, the findings of this kind of surveys will help to fine tune the health system to get maximum benefits of the investment. This will help the health planners to identify the institutions that need to be developed based on scientific evidence ensuring the availability and accessibility of quality health services.

I wish that the recommendations of this report will help to ensure high quality maternal and newborn health services throughout the country.

Maithripala Sirisena

Minister of Health,
Sri Lanka

Message from the Secretary to Ministry of Health



We are at a point in time when, the health service of the country is moving from a focus on coverage to quality and whole health system is gearing to implement the quality assurance mechanism at various levels. Maternal care services, one of the best services within the health system has already achieved universal coverage for most of evidence based interventions implemented through preventive and curative health services. Reflecting on the effective intervention coverage, Sri Lanka reports the best maternal mortality ratio, neonatal mortality rate and infant mortality rate in the region. Further, our maternal care programme has been recognized globally as being a highly effective system at a relatively low cost.

In order to sustain the already achieved satisfactory indicators and to ensure universal access to quality maternal and newborn services, consideration of the epidemiological data, fertility pattern, cost effectiveness, maximizing utilization of available resources, as well as the health seeking behavior and the expectation of the people has become an integral part of national, provincial and district level planning.

As such, in depth investigations on the current situation such as the EmONC needs assessment survey are needed to help the planning process. Therefore, I recommend that the findings of this document be used for policy making and planning at all levels. All stakeholders in maternal and neonatal health need to work together to maximize the use of the findings of the EmONC survey and ensure implementation of the recommendations of the report for the betterment of mothers and their newborns in this country.

Dr. Y.D. Nihal Jayathilaka

Secretary,
Ministry of Health,
Sri Lanka

Message from the Director General of Health Services



Despite Sri Lanka's achievements in the reduction of maternal mortality, neonatal mortality and infant mortality, there is much to be done in areas related to quality of service, management practices within hospitals, commitment of health staff, teamwork and leadership as well as attention given to client needs and their aspirations. Further reduction in maternal and neonatal mortality and morbidity will depend on addressing these issues. However, considering the current situation in Sri Lanka, universal answers will not always be cost effective, feasible or effective. As such, in depth and comprehensive analysis at institution, district and provincial level is necessary to help identify points for intervention.

This Emergency Obstetric and Neonatal Care Needs Assessment survey report has given a detailed description on availability, accessibility and utilization of maternal and neonatal health services together with the availability and distribution of human resources, drugs, equipment, infrastructure facilities and supportive services. This National report together with the Provincial reports would therefore be useful to identify deficiencies and inequities in distribution of resources and services and would contribute towards our efforts to further improve services for mothers in Sri Lanka and their newborn babies.

Dr. P.G. Mahipala,
Director General of Health Services
Ministry of Health,
Sri Lanka

FOREWORD

Improving the quality of maternal and newborn care services in the country is now receiving our attention with the view of further reducing the maternal and newborn mortality rates in Sri Lanka. As such, I see the Emergency Obstetric and Neonatal Care Needs Assessment Survey as a timely exercise.

In this study, we try to look at the facilities available for emergency obstetric and neonatal care and identify the gaps and deficiencies in existing services in government hospitals. It covers all the major hospitals providing comprehensive emergency obstetric and neonatal care services and a sample of divisional hospitals covering all the districts in the country.

I wish to take this opportunity to thank Sri Lanka College of Obstetrics and Gynecology and Sri Lanka College of Community Physicians for their valuable contribution.

I salute Dr Lalini Rajapaksa, the head of the research team and the other expert members of the team for coordinating and managing the complex tasks in a highly professional manner. I thank Dr. Nilmini Hemachandra and Dr Dammika Rowel, the two Consultant Community Physicians of the Family Health Bureau who are in charge of Maternal Care and Intranatal and Newborn Care respectively for spearheading this task.

Further, I hereby acknowledge with great appreciation the assistance extended by UNFPA, UNICEF and WHO, both in terms of their technical inputs and vital financial support.

This report also identifies the progress achieved since the last assessment which was done nearly thirteen years back, and I believe the findings of this study would be used as input for future planning and rational resource allocation.

Dr. Deepthi Perera

Director Maternal and Child Care
Family Health Bureau,
Ministry of Health
Colombo, Sri Lanka.

ACKNOWLEDGEMENTS

This report would not have been possible without the commitment and support from many individuals and organizations.

Dr. Ravindra Ruberu, former Secretary to Ministry of Health and Dr. Ajith Mendis, former Director General of Health Services provided invaluable guidance and administrative support to the initial activities of the EmONC needs assessment survey. Thereafter, Dr. Y.D. Nihal Jayathilake, the Secretary to Ministry of Health together with Dr. P.G. Mahipala, Director General of Health Services continued to provide leadership and guidance to this survey.

Dr. Terrance de Silva, former Deputy Director General (Medical Services) was a great strength during the initial stages of the assessment. The administrative support provided by Dr. Piyasena Samarakoon, former Acting DDG MS and the current DDG MS I, Dr. Lakshmi Somathunga is much appreciated. Dr. R.R.L.L.R. Siyambalagoda, Deputy Director General (Public Health Services) provided boundless support at all stages of the EmONC assessment and was a pillar of strength in resolving many challenges in the completion of the assessment.

The support and cooperation extended by Dr. Deepthi Perera, Director/MCH and Dr. Chithramalee de Silva Deputy Director/ MCH, particularly in resolving implementation and logistical issues throughout the assessment has been most valuable.

The key role played by the members of the National Core Group on the EmONC needs assessment survey in providing guidance at crucial stages of the assessment such as conceptualization, planning and implementation is greatly acknowledged.

The support extended by all Provincial Directors of Health Services and Regional Directors of Health Services in releasing Medical Officers of Health and providing them logistical support during this process has been of immense help to the successful completion of the assessment.

The dedication and commitment of Medical Officers of Maternal Child Health and Medical Officers Public Health in coordinating the assessment within the district is most appreciated, without which this assessment would not have been feasible.

The heads of institutions, matrons, sisters in charge, specialist obstetricians, specialist pediatricians, pharmacists, administrative and health staff in the selected institutions had been very responsive and most helpful during the data collection process. Through the diligence and commitment of the data collection teams comprising of medical officers from within the district (facility assessment) and institutions (prospective morbidity data collection) this assessment would not have been meaningful.

Appreciation is extended to Sri Lanka College of Obstetricians and Gynaecologists (SLCOG) and Sri Lanka College of Pediatricians (SLCP) for their unwavering support towards this assessment. Past presidents of SLCOG Dr. Sarath Amarasekera, Dr. Ananda Ranathunga and Dr. Hemantha Perera and the president of SLCOG Prof. Hemantha Senanayake are acknowledged for their technical expertise and productive comments.

Past presidents of SLCP Dr. Kalyani Guruge, Dr. Deepthi Samarage, and Prof. Ashvini Fernando and the president of SLCP Dr. Srilal de Silva are also acknowledged for their technical support at all times.

Development partners of the Ministry of Health, UNFPA, UNICEF and WHO provided their technical support and guidance with generous sponsorship all the stages of the assessment.

Deep appreciation and commendations are due to:

- Consultant pediatricians, Dr Medha Weerasekera and Dr Sandya Bandara made invaluable contributions by developing the newborn care modules for data collection.
- Specialist Obstetricians Dr Hemantha Perera and Dr Kapila Gunawardena gave their technical inputs to adapt the AMDD modules to suit the country context
- Dr. Kapila Jayaratne, Consultant Community Physician, Family Health Bureau and the in-charge of maternal mortality and morbidity surveillance, provided the maternal mortality data base for 2009 and 2010 for the analysis.
- Dr. Pamoda Madarasinghe, Dr. Duleepa Baranage and Dr. Indiwari Liyanage, research assistants for the EmONC needs assessment survey mixed their youth and creativity in all the stages of the assessment. Adaptation of the modules, maintaining the databases, data entry as well as data cleaning, coordinating with the hospitals, getting missing data made their life miserable, though they never complained.
- Dr. Shanika Senanayake and Dr. Samantha Jayasinghe, medical officers working in the maternal care unit of the Family Health Bureau. They were “the emergency team” that dealt with all the emergencies throughout the assessment.
- Dr. Nishamanie Karawita, former Health Officer, UNICEF was a great team player and provided valuable comments in improving the assessment. Her contribution in mobilizing funds is admirable.
- Dr. Anoma Jayathilake, National Professional Officer, WHO country office, Sri Lanka has been with the study team throughout and provided highly technical, balanced, politically and environmentally sound comments and constructive criticism throughout the assessment. Her support in crucial issues related to the assessment is commendable.
- Dr. Saramma Mathai, former programme coordinator at the Asia Pacific Regional office of UNFPA was instrumental in conducting the preliminary meetings and motivating the FHB, SLCOG, WHO, UNICEF and UNFPA to take up the EmONC assessment.

However, the more difficult path of implementation still lays ahead, the dedicated and active cooperation of all above and many more professionals would be vital for that journey.

CONTENTS

	Page
Dedication	iii
Emergency Obstetric and Neonatal Care Needs Assessment Research Team	iv
Core Group	v
Message from the Minister of Health	vii
Message from the Secretary to Ministry of Health	ix
Message from the Director General of Health Services	xi
Foreword	xiii
Acknowledgement	xv
Content	xvii
List of Tables	xxi
List of Figures	xxv
List of Abbreviations	xxvii
Chapter 1 Back ground	1
1.1 Objectives of the assessment	2
1.1.1 General Objectives	2
1.1.2 Specific Objectives	2
Chapter 2 Methodology	5
2.1.1 Sampling	5
2.1.2 Sampling for morbidity data collection and assessment of knowledge	7
2.2 Instruments for data collection	7
2.3 Pilot testing	8
2.4 Data collection	8
2.5 Data entry and analysis	8
2.5.1 Signal functions	9
2.5.2 Classification of health institutions by EmONC status	10
2.5.3 Indicators used in the study and their calculation	12
2.5.4 Other analysis	13
Chapter 3 Indicators of Emergency Obstetric Care	15
3.1 Selected characteristics of the districts and province	15
3.2 Provision of signal functions	18
3.3 Availability of EmONC services (Indicator 1)	21
3.4 Geographic distribution of EmONC facilities (indicator2)	23
3.5 Utilization of EmONC facilities	23

3.5.1	Proportion of births in EmONC facilities (Indicator 3)	23
3.5.2	Met need for EmONC Care (Indicator 4)	24
3.6	Caesarean Section (CS) as a proportion of all births (indicator 5)	24
3.7	Measures of quality of services	25
3.7.1	Direct obstetric case fertility rate (Indicator 6)	25
3.7.2	Intrapartum and very early neonatal death rate (Indicator 7)	26
3.7.3	Proportion of deaths due to indirect causes in EmONC facilities (Indicator 8)	26
3.8	Summary of EmONC indicators 1-8	27
3.9	New indicators developed in the current study	28
3.9.1	Refining indicator 2 (Geographic distribution of EmONC facilities)	28
3.9.1 (a)	Percentage area falling outside a defined buffer zone round CEmONC facilities	29
3.9.1.(b)	Percentage population falling outside a defined buffer zone round CEmONC facilities	30
3.10	Availability and accessibility	31
3.11	Composite indices developed in the study	32
3.11.1	Accessibility index for a specialist hospital (AISH)	32
3.11.2	Accessibility index for a specialist unit (AISU)	32
3.11.3	Accessibility index for a specialist Obstetrician (AISO)	32
3.12	Ministry of Health norms by district	33
Chapter 4	Selected Care Practices	37
4.1	Caesarean deliveries	37
4.1.1	Institutional Caesarean section rates	37
4.1.2	Indications for sections	38
4.1.3	Selected care practices for caesarean section	39
4.2	Use of partograph	40
4.3	Management of labour	41
4.3.1	Induction of labour	41
4.3.2	Selected practices related to labour	41
4.3.3	Referrals and average stay	42
4.4	Availability of guidelines and protocols for EmONC	42
Chapter 5	Maternal Mortality and Morbidity	45
5.1	Maternal mortality	45
5.2	Maternal morbidity	49
5.2.1	Admission Pattern	49
5.2.2	Delivery Pattern	49
5.2.3	Complications of pregnancy and labour	50
5.2.4	Co-morbidities	51
5.3	Transfers	52

Chapter 6	Resources for EmONC - Infrastructure	55
6.1	Facility profile	55
6.2	Labour room	56
6.3	Operating Theatre	58
6.4	Blood Transfusion services	58
6.5	Laboratory services	59
6.6	Facilities for infection control	60
6.7	Pharmacy and Medicines for EmONC services	60
6.8	Equipment Profile	63
6.9	Transport and Communication	64
6.10	Cost Incurred by patients by Institution	64
Chapter 7	Human resources for EmONC	67
7.1	Work load	67
7.1.1	Work load of different categories of car providers	67
7.1.1a	Variation among individual hospitals in different types of institutions	67
7.1.2	Work load for caesarean sections	69
7.2	In-service training profile	69
7.2.1	In -service Training profile of selected categories of personnel	70
7.2.2	In- service training profile of Medical officers	70
7.3	Knowledge among selected categories of care providers	71
7.3.1	Sample	71
7.3.2	Questionnaire	71
7.3.3	Analysis	72
7.3.4	Results	72
7.3.4.1	Institutional midwives	72
7.3.4.2	Nursing Officers	76
7.3.4.3	Medical Officers	80
Chapter 8	Preference for place of delivery	87
8.1	Preference for type of institutions	87
8.2	Movement of women between districts for delivery	89
Chapter 9	Institution based family planning services	95
Chapter 10	Neonatal Care services	99
10.1	Distribution of neonatal care facilities within the province	99
10.2	Availability and geographic distribution of neonatal care services	100
10.3	Morbidity profile	102

10.4	Interventions performed at the NICUs and SCBUs in the Western Province	104
10.5	Resources for neonatal services	105
10.5.1	Infrastructure	105
10.5.2	Human Resources	105
10.5.3	Guidelines, protocols, records and registers	107
10.5.4	Selected practices related to neonatal care Neonatal Examination	108
10.5.5	Drugs and equipment	109

Chapter 11	Recommendations	113
11.1	Signal Functions	113
11.2	Availability and accessibility	114
11.3	BEmONC facilities	114
11.4	CEmONC and 24*7 CEmONC facilities	119
11.5	Special Critical care facilities	120
11.6	Establishment of norms	120
11.7	Care practices	121
11.8	Quality Assurance process	122
11.9	Caesarean Sections	122
11.10	Maternal mortality and morbidity	122
11.11	Supportive services for EmONC	123
11.12	Training	123
11.13	Institutional based family planning services	123
11.14	Neonatal care services	124
11.14.1	Availability and accessibility	124
11.14.2	Establishment of Norms	125
11.14.3	Human Resource	126
11.14.4	Clinical standards and quality of care	126
11.14.5	Referral system	127
11.14.6	Neonatal transport system	127
11.14.7	Mother Baby Centres / Lactation Management Centres	127
11.15.	Others	127

LIST OF TABLES

Table 2.1	Hospitals included in the sample
Table 2.2	Indicators and their minimum acceptable levels
Table 3.1	Selected characteristics of the Western Province
Table 3.2	Classification at time of survey of hospitals included in the sample
Table 3.3	Classification of hospitals included in the sample by functional nature
Table 3.4	Distribution of births by district, type of hospital and by functional category of hospital
Table 3.5	Percentage of births in the sampled institutions (2011)
Table 3.6	Provision of signal functions in the 3 months preceding the survey by institution and district
Table 3.7	Availability facilities for provision of signal functions
Table 3.8	Institutions that have potential ability to provide signal functions by district
Table 3.9	Institutions that are able to provide CEmONC services on 24*7 by district
Table 3.10 (a)	Availability of EmONC facilities by district based on services provided during the 3 months prior to survey
Table 3.10 (b)	Availability of EmONC facilities by district based on the potential ability of institutions for providing signal functions
Table 3.10 (c)	Availability of institutions that are able to provide CEmONC services on a 24*7 basis by district
Table 3.11	Proportion of births occurring in CEmONC facilities by district and province
Table 3.12	'Met need' for CEmONC care by district and province
Table 3.13	Caesarean section as proportion of births registered in each district and province
Table 3.14	Direct obstetric case fatality rate by district and province
Table 3.15	Stillbirth rate per 1000 births by district and province
Table 3.16	Proportion of deaths due to indirect causes in CEmONC facilities
Table 3.17	Summary of EmONC indicators 1-8 by district
Table 3.18	Area per district falling outside a 30 km buffer zone from CEmONC and 24*7 CEmONC facilities
Table 3.19	Population without access to CEmONC and 24*7 CEmONC facility cover using a 30 Km buffer zone
Table 3.20	Specialist obstetricians per district according to Ministry of Health norms
Table 4.1	Caesarean section rates by institution, district and province
Table 4.2	Category of cesarean sections by institution and by districts based on one month prospective survey
Table 4.3	Indications for cesarean section
Table 4.4	Service provision for caesarean sections

Table 4.5	Use of partograph in institutions
Table 4.6	Induction of labour in specialist and non-specialist institutions in Western Province
Table 4.7	Practices related to labour
Table 4.8	Referral practices and duration of stay by institution and district
Table 4.9	Availability of guidelines and protocols by institution
Table 5.1 (a)	Causes of maternal mortality by district 2001-2010 Colombo
Table 5.1 (b)	Causes of maternal mortality by district 2001-2010 Gampaha
Table 5.1 (c)	Causes of maternal mortality by district 2001-2010 Kalutara
Table 5.2	Admission patterns by type of institution
Table 5.3	Delivery patterns in specialist and non- specialist institutions in the Western province
Table 5.4	Complications of pregnancy and labour in specialist and non- specialist institutions in Western province
Table 5.5	Co-morbidities in specialist institutions and non- specialist institutions in Western province
Table 5.6	Pattern of transfers in the Western Province
Table 5.7	Pattern of morbidity in transferred patients in Western Province
Table 6.1	Summary of services provided by institution
Table 6.2	Summary of institutional profiles in each district
Table 6.3 (a)	Summary of labour room facilities by institution - infrastructure
Table 6.3 (b)	Summary of labour room facilities by institution - infrastructure
Table 6.3 (c)	Summary of labour room facilities by institution - infrastructure
Table 6.4	Operation Theatre facilities by institution
Table 6.5	Blood transfusion services by institution
Table 6.6	Laboratory services by Institution
Table 6.7 (a)	Availability of Infection control measures by institution
Table 6.7 (b)	Infection control measures by institution
Table 6.8 (a)	Availability of essential drugs by institution
Table 6.8(b)	Availability of essential drugs by institution
Table 6.9	Pharmacy management practices by institution
Table 6.10	Average daily use of selected equipment by institution
Table 6.11	Average daily use of theatre facilities for obstetric care by institution
Table 6.12	Transport and communication facilities by institution
Table 6.13	Costs incurred by patients by institution
Table 7.1	Work load of selected categories of care providers for normal deliveries by institution
Table 7.2	Workload for caesarean sections by institution
Table 7.3	In-service training profile of selected categories of personnel in the province
Table 7.4	In-service training profile of medical officers
Table 7.5	The number of respondents in each category

Table 7.6	Causes given by midwives for each condition
Table 7.7	Frequency distribution of the causes of PPH as given by midwives
Table 7.8	Frequency distribution of the organs affected by severe PIH (Responses from midwives)
Table 7.9	Frequency distribution of the causes for sudden collapse after delivery (Responses from midwives)
Table 7.10	Frequency distribution of the Symptoms of severe heart disease (Responses from midwives)
Table 7.11	Marks obtained by the midwives for each component
Table 7.12	Summary of marks obtained by the midwives
Table 7.13	Causes given by nursing officers for each condition
Table 7.14	Frequency distribution of the causes of PPH (Responses by nursing officers)
Table 7.15	Frequency distribution of the organs affected by severe PIH (Responses from nursing officers)
Table 7.16	Frequency distribution of the causes for sudden collapse after delivery (Responses from nursing officers)
Table 7.17	Frequency distribution of the symptoms of severe heart disease (Responses from nursing officers)
Table 7.18	Marks obtained by the nursing officers for each component
Table 7.19	Summary of marks obtained by the nursing officers
Table 7.20	Causes given by medical officers for each condition
Table 7.21	Frequency distribution of the causes of PPH (Responses by medical officers)
Table 7.22	Frequency distribution of the organs affected by severe PIH (Responses from medical officers)
Table 7.23	Frequency distribution of the causes for sudden collapse after delivery
Table 7.24	Frequency distribution of the Symptoms of severe heart disease (Responses from medical officers)
Table 7.25	Marks obtained by the medical officers for each component
Table 7.26	Summary of marks obtained by the medical officers
Table 8.1	Calculation of percentage completeness of the data reported from the field
Table 8.2	Place of delivery reported in field data
Table 8.3	Deliveries occurring in government hospitals by district
Table 8.4	Deliveries occurring within the district, going out and coming in to the district
Table 9.1	Institution based family planning services in Western Province
Table 9.2	Institution based family planning clinics in Western Province
Table 9.3	Logistics and documentation of the Family Planning Services
Table 10.1	Facilities available for neonatal care in the Western Province
Table 10.2	Availability of NICU and SCBU by population and extent
Table 10.3	Morbidity pattern at the Institutions with NICU and SCBU in the Western Province
Table 10.4	Interventions performed at the NICU and SCBUs
Table 10.5(a)	Infrastructure for neonatal services by institution and district

Table 10.5(b)	Infrastructure for neonatal services by institution and district
Table 10.6	Human resource distribution
Table 10.7	Availability of protocols and guidelines at NICU/SCBU
Table 10.8	Registers and records available at NICU/ SCBU
Table 10.9	Selected practices related to neonatal care by institution and district
Table 10.10	Drugs available at NICU/ SCBU
Table 10.11	Facilities for infection control by institution and district
Table 10.12	Equipment available at NICU/ SCBU
Table 10.13	Drugs and equipment available for neonatal resuscitation at NICU/ SCBU

LIST OF FIGURES

Figure 1	Government medical Institutions in the Western Province
Figure 3.1	Equity Assessment Matrix of the 8 EmONC indicators
Figure 3.2	Facilities capable of providing CEmONC and 24*7 CEmONC care
Figure 3.3	Scatter plot of CEmONC facilities by indicators one and two
Figure 3.4	Scatter plot of 24*7 CEmONC facilities by indicators one and two
Figure 3.5	Summary of indicators 1 and 2, new indicators and utilization indicators by district
Figure 5.1	Trends in maternal mortality by district 2001-2010
Figure 5.2	Pattern of obstetric transfers in the Western province
Figure 8.1	Distribution of births in the government institution by type of hospital and district -2011
Figure 8.2	Distribution of births in the government hospitals by functional classification and district -2011
Figure 8.3	Distribution of place of delivery and within district migration for delivery
Figure 10.1	Pattern of neonatal transfers in the Western Province

LIST OF ABBREVIATION

24*7	24 hours a day, every day of a week
AISH	Accessibility Index for a Specialist Hospitals
AISU	Accessibility Index for a Specialist Unit
AISO	Accessibility Index for a Specialist Obstetrician
AMBU	Artificial Manual Breathing Unit
AMDD	Averting Maternal Death and Disability Program
BB	Blood Bank
BEmONC	Basic Emergency Obstetric and Neonatal Care
BH	Base Hospital
BHT	Bed Head Tickets
BT/CT	Bleeding Time/ Clotting Time
CEmONC	Comprehensive Emergency Obstetric and Neonatal Care
CS	Cesarean Section
CSSD	Central Sterile Supply Department
CTG	Cardiotocography
DGH	District General Hospital
DH	Divisional Hospital
DMPA	Depot Medroxy Progesterone Acetate
ECG	Electrocardiography
ECP	Emergency Contraceptive Pills
EmOC	Emergency Obstetric Care
EmONC	Emergency Obstetric and Neonatal Care
FFP	Fresh Frozen Plasma
FHB	Family Health Bureau
FP	Family Planning
H 830	Monthly Maternity Statistics
HO	House Officer
ICU	Intensive Care Unit
IUD	Intra Uterine Device
IUGR	Intra Uterine Growth Retardation
IV	Intra venous
LMC	Lactation Management Centre
LR	Labour room
LRT	Ligation and Resection of Tubes
LSCS	Lower Segment Cesarean Section
MBC	Mother Baby Centre
MgSO ₄	Magnesium Sulphate
MI	Myocardial Infarction

MLT	Medical Laboratory Technician
MO	Medical Officer
MOH	Medical Officer of Health
MOMCH	Medical Officer/ Maternal and Child Health
ND	Normal Delivery
NICU	Neonatal Intensive Care Unit
NO	Nursing Officer
NR	Neonatal Resuscitation
OT	Operation Theatre
PCV	Pack Cell Volume
PGH	Provincial General Hospital
PHM	Public Health Midwife
PHO	Paediatric House Officer
PIH	Pregnancy Induced Hypertension
PPH	Post Partum Haemorrhage
PNS	Post Natal Side
RDHS	Regional Director of Health Services
RG	Register General
RMSD	Regional Medical Supplies Division
SCBU	Special Care Baby Unit
SHO	Senior House Officer
SLCOG	Sri Lanka College of Obstetricians and Gynaecologists
SLCP	Sri Lanka College of Paediatrics
TH	Teaching Hospital
VOG	Visiting Obstetrician and Gynaecologist
UNFPA	United Nations Population Fund
UNICEF	United Nations Children's Fund
WHO	World Health Organization

Chapter 1

Background

Sri Lanka has achieved much in terms of reducing maternal and infant mortality. The decline in mortality, trends in cause specific mortality and the factors contributing to these have been well documented. However, these favourable national statistics hide regional and sectoral differences. The differentials in maternal mortality have narrowed over time but are still observed.

It is well known that risk factors while identifying groups of women who may be at increased risk of a complication does not necessarily predict an individual woman who is likely to develop a complication. These complications can be treated successfully if high quality, emergency obstetric care is available and accessible to the women who need them. Further improvements in maternal and infant mortality in Sri Lanka will largely depend on the capacity of the health system to respond to obstetric and new born emergencies and complications. Thus, the strengthening of Emergency Obstetric and Newborn Care (EmONC) services and reducing disparities in availability and accessibility of these services are of importance.

Historically, the status of maternal care has been tracked mainly using a health impact indicator, the maternal mortality ratio. Outcome indicators such as use of antenatal care, coverage of immunisation with tetanus toxoid and skilled attendance at birth have been added on. However, these outcome indicators do not capture the quality of provision of care in totality and do not align well with the impact

indicator used. In more recent times, indicators of health systems outputs or process indicators have been promoted and there has been publications such as the “UN guidelines for monitoring the availability and use of obstetric services” (1997) and “Monitoring Emergency Obstetric Care: A handbook” (WHO 2009). These indicators have the advantage of being appropriate for use both nationally and regionally, but In Sri Lanka, these have been under used in routine monitoring of maternal care up to date. With decreasing maternal mortality and near universal coverage of antenatal care, immunisation and skilled attendance at birth, these traditional indicators alone have become inadequate to monitor short term progress in care and to capture changes that are necessary to further fine tune the interventions and services. In this scenario, national and regional EmONC care indicators have an important role to play in the monitoring of maternal care provision in the country. Such an assessment was last carried out nearly 10 years ago.

Therefore, the current EmONC needs assessment was planned to examine the ability of the different grades of health institutions at regional and national level to provide necessary life-saving care to pregnant women and their newborn and also to serve as a benchmark against which future progress can be measured.

This report presents the data for the Western province collected as part of the national survey. The findings would be useful in improving the services in the province.

1.1 Objectives of the assessment

The objectives of the assessment were as follows:

1.1.1 General objective

Describe the current availability, geographical distribution, level of utilization, selected aspects of quality of EmONC services in the Western Province.

1.1.2 Specific objectives

To describe by district and province;

1. The availability and geographic distribution of comprehensive and basic Emergency Obstetric care facilities in terms of provision of signal functions and indicators of EmONC

2. Selected care practices
3. Maternal mortality and morbidity patterns
4. Resources for emergency obstetric care i.e.
 - Infrastructure, equipment and essential drugs
 - Human resources and training
5. Describe institution based family planning services
6. Neonatal care services and morbidity pattern

Chapter 2

Methodology

The Western Province of the country consists of three districts: Colombo, Gampaha and Kalutara. It is bounded by the sea on the West, the Southern Province to the South, Sabaragamuwa to the East, and the North Western Province to the North. The administrative and commercial capital of the country lies in the Colombo District and the province is considered the most resource rich in the country. The province also has the highest population density.

2.1 Sampling

The EmONC assessment was carried out in all districts. Within a district all hospitals where


specialist obstetric services are expected to be available were included in the sample; (i.e. base hospitals and above). Other institutions where deliveries occur but where specialist services are not available were sampled. The number of institutions below the level of a base hospital sampled per district depended on the percentage of deliveries performed in these institutions out of the total in the district. The required number of smaller institutions per district was selected probability proportional to the number of deliveries in each institution.

All government health facilities in the Western province are shown in figure 01 and those selected in to the sample in table 2.1.

Table 2.1: Hospitals included in the sample

Type of facility	Colombo	Gampaha	Kalutara
TH/PGH/DGH	TH - Castle Street TH - De Soysa TH - Sri Jayawardhanapura TH -Colombo South	TH -Colombo North DGH - Gampaha DGH - Negombo	DGH - Kalutara
BH	BH - Avissawella BH- Homagama BH - Mulleriyawa	BH - Watupitiwala	BH - Panadura BH - Horana
Divisional Hospital	DH - Piliyandala	DH - Mirigama DH - Kiribathgoda DH - Minuwangoda	DH - Pimbura DH - Matugama DH - Beruwala

Figure 1: Government medical institutions in the Western Province



2.1.2 Sampling for morbidity data collection and assessment of knowledge

Morbidity data were collected prospectively for a period of 4 weeks from all obstetric units and from the NICUs and SCBUs in the selected institutions.

Knowledge was assessed among three categories of care providers i.e., midwives, nursing officers in obstetric units, labour rooms and the first contact medical officers in the units (Intern House Officers / Medical Officers). The data collection tool was administered to all staff members in the above categories who were present on the day the assessment teams visited a given unit.

2.2 Instruments for data collection

It was decided that the generic modules developed by the Averting Maternal Death and Disability Program (AMDD) at Columbia University and its partners (including UNICEF, UNFPA, WHO, Care and Save the Children) be used for the assessment. This set of documents to EmONC services, the added “N” emphasizing the fact that information on emergency services for new-borns as well as mothers is included. This report therefore uses the term EmONC.

This set of instruments have been developed based on the experiences gained through the conduct of facility-based assessments of the availability, utilization and quality of emergency obstetric and newborn care (EmONC) in over 50 countries in Africa, Asia and Latin America. The modules use various data collection methods including interviews, extraction of information (data) from records and registers, and observation.

These modules were adapted to suit the Sri Lankan context. Instruments that were available locally for evaluation of obstetric practice were also examined and used where appropriate. The AMDD generic module on knowledge was not used. A special module was developed for this purpose which specifically addressed applied knowledge required to deal with common obstetric emergencies.

A neonatal module was developed to help identify current availability as well as needs. Neonatal care services in the country are in the process of being organised. The information collected has been planned with a view to providing inputs in to this process.

The adapted tools were discussed in detail with a wider group of stake holders consisting of technical experts from the College of Obstetricians and Gynaecologists (SLCOG), College of Paediatricians (SLCP), Perinatal Society of Sri Lanka, specialists in the Family Health Bureau, representatives from the Ministry of Health and Provincial Health Administration. Modifications were made during the consultative process and consensus reached. The core group approved the finalized set of tools.

The list of data collection modules used in the survey is given below:

- Module 1: Identification of facility and infrastructure: covers background information on the facility - including size/capacity, overall infrastructure, transport, communication
- Module 2: Human resources: signal functions the staff provide, staffing situation 24 hours a day 7 days a week
- Module 3: Essential drugs, equipment & supplies
- Module 4: Facility case summary: these data include deliveries, obstetric complications, caesarean deliveries, maternal deaths, intrapartum stillbirths
- Module 5: EmONC signal functions & other important services
- Module 6: Partograph review
- Module 7: Assessment of health provider knowledge

- Module 8: Review of 10 BHTs per unit where delivery was by caesarean section

In addition to the above modules the following tools were developed specifically for the current survey:

- A module on institution based family planning services,
- Checklist for assessment of labour rooms, post natal wards, SCBU/NICU, a tool for collection of morbidity data from bed head tickets,
- Formats for qualitative data collection on the management of the last averted maternal death and a severe neonatal morbidity,
- Formats to obtain comments from the specialist obstetricians and specialist paediatricians / neonatologists on how to improve the current maternal mortality review process and the perinatal mortality conference,
- Format for collection of data on place of delivery from each PHM area for a specified period of 3 months (quarter of the year).

The data collector's manual provided by AMDD was modified to suit the local survey.

Since there is a well-established national maternal death review process in place, no attempt was made to collect information on maternal deaths during field data collection. Data from the Maternal Mortality Data Base maintained in the FHB was used for analysis. Deaths that had occurred in 2009 and 2010 were analyzed in detail.

The methodology of the survey is described in detail in the national report.

2.3 Pilot test

The pilot test of survey procedures was carried out in the Kalutara district. All institutions included in the sample for the district were included in the pilot test. The pilot test also examined the feasibility of

obtaining morbidity data from registers, prospectively from admissions and retrospectively from discharge BHTs and prospectively from discharge BHTs. It was seen that the most feasible method for collection of morbidity data was prospectively from discharges and the best method of getting an accurate diagnosis and information on co-morbidities was to entrust the task to a medical officer in the ward.

2.4 Data collection

In a given district, the Medical Officer of Maternal and Child Health (MOMCH) was responsible for collecting the necessary data from all institutions included in the sample. The Regional Director of Health Services (RDHS) facilitated the process by arranging and authorising transport for the teams of data collectors and authorising duty leave for data collection.

Two to three data collection teams were selected per district depending on the number and size of institutions. Each team comprise of 3 persons selected from among senior Medical Officers of Health (MOH), MO/public health and MO/planning. Persons with experience in Obstetrics and Paediatrics were recruited as data collectors and all data collectors were trained in the use of the modules.

Morbidity data was collected from all institutions prospectively for one month. All discharges from a unit over a period of 4 weeks from the date of commencement were collected. This was carried out by a medical officer attached to the unit and designated by the specialist obstetrician or paediatrician for this purpose. The nursing sister in charge of the ward was responsible for ensuring that all discharges were included in the forms and was given a specially designed check list for the purpose.

2.5. Data entry and analysis

All data entry forms were checked for completeness and an attempt was made to obtain any missing data from the districts/ the classification of health institutions concerned. Data were entered using Epi data and Excel formats. Ten per cent of all entries were cross checked by a second person.

Institutions published by the Ministry of Health were used in the analysis (Ministry of Health 2010). At the time of data collection it was noted that some of the institutions had been upgraded from the classification used in the sampling. In addition to this a functional classification based on availability of specialist services was also used.

The data were analysed according to the EmONC indicators described in the manual “Monitoring Emergency Obstetric Care: A handbook” (WHO 2009).

2.5.1 Signal functions

The term Emergency Obstetric and Neonatal Care refers to the provision of a list of life saving services or signal functions which defines a health facility in respect of its ability to treat obstetric and newborn emergencies. Signal functions used to classify a facility are given below (Box 2.1).

Box 2.1: List of signal functions

EmONC Signal Functions
1. Administer parenteral antibiotics
2. Administer uterotonic drugs (i.e. parenteral oxytocin)
3. Administer parenteral anticonvulsants for pre eclampsia and eclampsia (i.e. magnesium Sulphate)
4. Manually remove the placenta
5. Remove retained products
6. Perform assisted vaginal delivery (e.g. vacuum extraction, forceps delivery)
7. Perform basic neonatal resuscitation (e.g. with bag and mask)
8. Perform surgery (e.g. caesarean section)
9. Perform blood transfusion

2.5.2 Classification of health institutions by EmONC status

A basic emergency obstetric care facility (BEmONC) is one in which functions 1 – 7 are performed while a comprehensive obstetric care facility (CEmONC) is one in which all functions 1 – 9 are performed .

In Sri Lanka, removal of retained products (signal function 5) is not recommended in institutions where there are no specialist obstetric services. Institutions that provide signal functions 1-7 excluding signal function 5 are referred to in the current analysis as a modified BEmONC facility.

The first EmOC needs assessment carried out in the year 2000, examined the provision of the first 4 basic signal functions in addition. The present study this was also examined for purposes of comparison.

According to the reference “Monitoring Emergency Obstetric Care: A handbook” WHO (2009) an institution is classified as BEmONC or CEmONC based on providing the signal functions in the three months preceding the assessment. Data collected through direct inquiry is used for this classification.

It is noted that there may be many reasons for not performing a function in the stipulated period such as policy decisions on service provision, (For example in Sri Lanka, removal of retained products of conception is performed only in institutions where there are specialists in obstetrics) absence of cases needing specific services, lack of persons, skills or infrastructure necessary for the performance of a particular function. Therefore, the present analysis used a set of objective criteria that was agreed upon by the core group as necessary for the provision of each signal function to identify institutions that are “potentially able to provide signal functions”. The set of criteria are given in Box 2.2.

Box 2.2: Criteria considered as necessary for the provision of signal functions

1. Drugs
 - a. at least one parenteral antibiotic
 - b. at least one parenteral uterotonic drug
 - c. Magnesium sulphate /parenteral anticonvulsants
 - d. Pethidine
 - e. Adrenaline
2. Equipment (in working order)
 - a. Vacuum extractor / forceps
 - b. Neonatal AMBU bag (Artificial Manual Breathing Unit)
3. Personnel
 - a. At least one specialist obstetrician
 - b. At least one MO blood bank
 - c. At least one MO trained in anaesthesia
 - d. House officers / Medical officers
 - e. Nursing officers
 - f. Midwives
4. Infrastructure
 - a. Electricity
 - b. Continuous water supply
 - c. Functional operating theatre
 - d. Functional blood bank
5. Training issues not being given as a reason for not providing the following services in data collection format 5B
 - a. Manual removal of placenta
 - b. Assisted vaginal deliveries
 - c. Neonatal resuscitation

The conditions were used to categorise the institutions on their potential ability to perform signal functions 1-9. The institutions that fulfilled the criteria given box 2.2 are referred to in the analysis as being “potentially able to provide the signal functions”.

Emergency functions should be available for patients on a 24*7 basis i.e. throughout the day every day of the week. Therefore, the ability of institutions to

provide services 24*7 were examined based on the set of minimum criteria given below (Box 2.3). The emphasis was on the ability to provide surgical, anaesthetic and blood transfusion services on a 24*7 basis. The institutions that fulfilled the criteria are referred to as “institutions able to provide CEmONC functions on a 24*7 basis” in the analysis. This analysis was confined to institutions that had the ability to provide CEmONC services.

Box 2.3: Minimum criteria used in classifying an institution as capable of providing signal functions 24*7

1. Administration of parenteral antibiotics
 - a. availability of parenteral antibiotics
 - b. trained staff capable of administering parenteral antibiotics: medical officers and nursing officers with basic training
2. Administration of uterotonic drugs
 - a. availability of uterotonic drugs
 - b. trained staff capable of administering uterotonic drugs: medical officers and nursing officers with basic training
3. Administration of parenteral anticonvulsants
 - a. availability of parenteral anticonvulsants
 - i. Availability of Magnesium Sulphate
 - b. trained staff capable of administering parenteral anticonvulsants: medical officers and nursing officers with basic training
4. Performing manual removal of placenta
 - a. availability of pethidine
 - b. trained staff capable of performing manual removal
 - i. medical officers with basic qualifications
 - ii. training issues not mentioned as reason for not performing the activity in data collection format 5B
5. Performing removal of retained products:
 - a. this function is recommended to be performed where there is a specialist obstetrician present in Sri Lanka
6. Performing assisted vaginal deliveries
 - a. Availability of vacuum extractors or forceps
 - b. Trained staff capable of performing assisted vaginal deliveries
 - i. medical officers with basic qualifications
 - ii. training issues not mentioned as reason for not performing the activity in format 5B
7. Performing basic neonatal resuscitation (NR)
 - a. availability of a neonatal resuscitation area in the LR
 - b. Availability of neonatal AMBU bag
 - c. trained staff capable of performing neonatal resuscitation
 - i. nursing staff in LR / PNS trained on NR
 - ii. OR mid wife trained on NR
 - iii. OR medical staff trained on NR
8. Performing emergency obstetric surgeries
 - a. Functional / operational operating theatre
 - b. Electricity
 - c. Functional generator
 - d. continuous source of water
 - e. At least two specialist obstetricians
 - f. Two or more persons trained in anesthesia MO or specialist
9. Performing blood transfusions
 - a. Functional blood bank
 - b. At least three MOs Blood Transfusion service
 - c. Electricity
 - d. Functional generator (absolute necessity)
 - e. continuous source of water

In addition to the above the following facilities were considered as essential in the provision of continuous EmONC services.

1. 24*7 availability of electricity
 - a. Primary source of electricity to be obtained from the national grid
 - b. Continuous electricity supply
 - c. Electricity to be available at time of interview
 - d. Functional backup generator to be available
 - e. Days without electricity during the past month to be 'zero'
2. 24*7 availability of water supply
 - a. Primary source of water to be obtained from piped water
 - b. Continuous water supply
 - c. Adequacy of the water supply
 - d. Days without water during the past month to be 'zero'

3. 24*7 availability of communication facilities
 - a. Availability of land phones in the facility
 - b. Communication facility to be available 24*7

The analysis thus described each institution under 3 scenarios of provision of signal functions namely:

- Provided the function in the three months prior to survey,
- Potentially able to provide the function
- Are able to provide the functions on a 24*7 basis.

2.5.3 Indicators used in the study and their calculation

The indicators used and their minimum acceptable level are given in table 2.1. The indicators were defined and calculated as per guidelines given in the manual quoted above and the details of calculation are given in Annex I.

Table 2.1 Indicators and their minimum acceptable levels

Indicator	Minimum acceptable level
Availability of EmONC	Five EmONC facilities per 500000 population out of which at least one should provide comprehensive care.
Geographic distribution of EmONC facilities	No standard set
Proportion of all births in Basic and Comprehensive EmONC facilities	No standard set Long term objective is 100% of births to take place in a facility that can deal with emergencies.
"Met need" for EmONC	Goal is that 100% of women with a complication will receive EmONC
Caesarean sections as a % of all births	5-15% is the recommended rate in the WHO hand book in 1985
Direct obstetric case fatality rate	Maximum acceptable level less than 1%
Intrapartum and very early neonatal death rate	No standard set
Proportion of deaths due to indirect causes in EmONC facilities	No standard set

Indicators one and two were calculated based on all three scenarios of provision of signal functions given above. The study used a sample of 46 institutions below the level of a base hospital. The institutions sampled accounted for nearly 45% of births in this type of health facility. The findings from the sample are extrapolated to estimate the number of institutions below the level of a Base Hospital that may be classified as having provided modified BEmONC services in the three months prior to survey and institutions that have the potential ability to provide modified BEmONC functions. Reasons for not being able to provide each signal functions were examined. The classification of institutions on the ability to provide services on a 24*7 basis is confined to institutions that are able to provide CEmONC services.

The district population published by the Department of Census and Statistics based on the census 2012 was used for indicators which required the total population as the denominator (Date of census is given as the 20th March 2012). The number of births registered for the year 2011 obtained from the Registrar General by “place of occurrence of birth” was used when indicators required the number of births per district as the denominator. The number of births by “place of occurrence” was preferred over the number of births by “place of usual residence” since the indicators examine the services provided by the institutions in a district. Where appropriate the number of deliveries occurring in government hospitals obtained from the RDHS is used. The number of maternal deaths per district and causes of death for the year 2009 & 2010 were obtained from the Family Health Bureau as this was the last two years for which maternal death reviews have been completed.

2.5.4 Other analysis

The morbidity rates were estimated based on the sample by level of institution; namely TH/PGH, DGH, BH with specialist obstetricians, BH upgraded but without specialists, institutions below the level of BHs. National morbidity rates are calculated based on weighted averages.

Resources available for provision of EmONC services are described in terms of infrastructure, human resources, drugs and equipment.

A set of indicators similar to what is described for Emergency obstetric care is not available for Neonatal care services. The morbidity pattern, infrastructure, human resources, drugs and equipment available are described by type of institution in the provincial reports.

The response to the attempt to collect qualitative data on a near miss and improvements to the current maternal mortality and perinatal mortality investigation procedures was not successful. This data is not presented in the report due to the high non response rate.

A provincial analysis based on the broad framework agreed upon by the core group was presented and discussed in two provinces, one with teaching hospitals and the other without. The objective was to obtain feedback on how best to present the data collected so that it would be useful for provincial administrators and clinicians. The format of the final provincial reports and the national report were discussed at the core group and consensus obtained.

Chapter 3

Indicators of Emergency Obstetric Care

3.1 Selected characteristics of the Western Province

Selected characteristics of the Western province are described in table 3.1. It is noted that the total deliveries in government institutions based on the hospital statistics from the office of the Regional Director of Health Services (RDHS) reported for the year 2011, is much less than the births reported by the Registrar General (RG) in all 3 districts. Note that the RGs data is reported by place of delivery. RGs data records an excess of 9637, 3402, 231 births in Colombo, Gampaha and Kalutara respectively.

Table 3.2 gives the classification of hospitals by type of facility. It is noted that some institutions have been upgraded since the selection of the sample. However, specialist services were not available in some of the institutions upgraded to BH, at the time of survey. The new terminology is used in the presentation of results.

It must be noted that although Sri Jayawardanapura hospital was included in the sample, data from this institution was not available for analysis.

Table 3.1: Selected characteristics of the Western Province

Characteristic	Colombo	Gampaha	Kalutara	Western Province
Population in 2012	2,323,826	2,298,588	1,214,880	5,837,294
Geographic extent in km ²	699	1387	1598	3684
Total deliveries in government hospitals in 2011 (H830)	45251	24055	16716	86022
Total births In the district in 2011 (RG)	54,888	27,457	16,947	99292

Table 3.2: Classification of hospitals included in the sample at the time of survey

Type of facility	Colombo	Gampaha	Kalutara
TH/PGH/DGH	TH - Castle Street TH - De Soysa TH - Sri Jayawardhanapura TH -Colombo South	TH -Colombo North DGH - Gampaha DGH - Negombo	DGH - Kalutara
BH	BH - Avissawella BH - Homagama BH - Mulleriyawa	BH - Watupitiwala BH - Mirigama BH - Kiribathgoda	BH - Panadura BH - Horana BH - Pimbura
Divisional Hospitals	DH - Piliyandala	DH - Minuwangoda	DH - Matugama DH - Beruwala

Table 3.3 describes the functional classification used in the analysis and the colour coding used in the table to distinguish between the categories is carried throughout the analysis.

Table 3.3: Classification of hospitals included in the sample by functional nature

<i>Functional classification</i>	<i>Colombo</i>	<i>Gampaha</i>	<i>Kalutara</i>
Hospitals with specialized units			
More than one Obstetric unit	TH Castle Street TH De Soyza TH -Colombo South BH Awissawella BH Homagama	TH- Colombo North DGH -Gampaha DGH -Negambo BH -Wathupitiwala	DGH -Kaluthara BH -Panadura BH -Horana
One unit with two specialist obstetricians			
One unit with one specialist obstetrician	BH Mulleriyawa		
Upgraded hospitals without a specialist obstetrician at the time of the survey		BH Meerigama BH Kiribathgoda	BH Pimbura
Divisional Level/ non specialist units	DH - Piliyandala	DH - Minuwangoda	DH - Matugama DH - Beruwala

Table 3.4 shows the distribution of births in the province based on the type of facilities and by the functional classification.

Table 3.4: Distribution of births by district, type of hospital and by functional category of hospital

	Colombo		Gampaha		Kalutara		Western Province	
Total reported deliveries in government institutions in 2011 (H830)	45251		24055		16716		86022	
Based on the type of facility								
TH / PGH / DGH	37329	82.5	20046	83.3	8708	52.1	66083	76.8
BH	7184	15.9	3888	16.2	7709	46.1	18781	21.8
Divisional Hospital	738	1.6	121	0.5	299	1.8	1158	1.4
Based on the functional category of hospital								
More than one specialized unit	44271	97.8	23920	99.4	16407	98.2	84598	98.3
One unit with two specialist obstetricians	0	0.0	0	0.0	0	0.0	0	0.0
One unit with one specialist obstetrician	242	0.5	0	0.0	0	0.0	242	0.3
Upgraded units with no specialist obstetrician at time of survey	0	0.0	14	0.1	10	0.1	24	0.02
Divisional level non specialized hospital	738	1.7	121	0.5	299	1.8	1158	1.4

Table 3.4 shows that less than 1.5% of deliveries in the government institutions in the province take place in institutions where specialist services are not available. This proportion is 1.7%, 0.6%, and 1.9% in the districts of Colombo, Gampaha and Kalutara respectively.

Table 3.5 shows that the institutions included in the study sample accounts for 98.9% of the births in the province. However it must be noted that Sri Jayawardhanapura hospital did not provide data for this analysis thus the institutions for which data is available account for 91.29% of births in the Colombo district and 89.8% in the province.

3.5: Percentage of Births in the sampled institutions (2011)

	COLOMBO			GAMPAHA			KALUTARA			WESTERN PROVINCE		
	Specialized institutions	Non specialist Institutions	District	Specialized institutions	Non specialist Institutions	District	Specialized institutions	Non specialist Institutions	District	Specialized institutions	Non specialist Institutions	Province
No. of institutions in the district	7	11	18	4	24	28	3	18	21	14	53	67
No. of births in 2011	44513	738	45251	23920	135	24055	16407	309	16716	84840	1182	86022
No. of institutions in the sample	6	1	7	4	3	7	3	3	6	14	7	20
No. of births in 2011 in the sampled institutions	40636	14	40650	23920	40	23960	16407	145	16552	80963	199	77285
Percentage coverage of births by the sample	91.29	1.9	89.8	100	29.6	99.6	100	46.9	99.0	100	16.8	89.8

3.2 Provision of signal functions

A list of functions that can save lives of women in an obstetric emergency called 'signal functions' have been identified by WHO and partners. The provision of these functions during the three months prior to the survey is used to classify institutions as having basic or comprehensive emergency obstetric and neonatal care functions.

Table 3.6 describes the provision of signal functions in the sampled institutions in the province. It is noted that in Sri Lanka, removal of retained products is not recommended in institutions where there are no specialist obstetric services.

The first EmONC needs assessment carried out in the year 2000, examined the provision of the first 4 basic signal functions. Therefore in the present study too, this was examined for purposes of comparison.

The table (3.6) shows that excluding SJP the district of Colombo has 4 institutions that have provided all 9 signal functions and can be classified as CEmONC facilities. In Gampaha and Kalutara there are 4 and 3 institutions respectively that have provided comprehensive emergency care in the 3 months preceding the survey.

In addition to the above there are two other institutions in the Colombo district that have provided all signal functions except neonatal resuscitation.

Further to the above, the availability of facilities for the provision of signal functions was examined and is given in table 3.7. Based on this table, the institutions that are capable of providing the signal functions are described in table 3.8.

Table 3.6: Provision of signal functions by institution and district (during the last 3 months)

Hospital	Colombo							Gampaha							Kaluthara						
	TH CSHW	TH DMH	TH Colombo South	BH Avissawella	BH Homagama	BH Mulleriyawa	DH Piliyandala	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	BH Mirigama	BH Kiribathgoda	DH Minuwangoda	DGH Kaluthara	BH Panadura	BH Horana	BH Pimbura	DH Mathugama	DH Beruwala	
Administering of parenteral antibiotics							0						0						0		
Administering of parenteral oxytocins													0	0							
Administering of parenteral anti convulsants							0					0	0	0				0	0	0	
Performing manual removal of placenta							0					0	0						0	0	
Performing assisted vaginal delivery							0						0	0					0	0	
Removal of retained products of conception													0								
Performing neonatal resuscitation					0							0	0	0				0	0	0	
Performing obstetric surgeries													0								
Performing blood transfusions													0								
		Signal function performed during the last three months																			
		Signal function not performed during the last three months																			
0		Signal function not performed due to absence of cases needing the service																			

Table 3.7: Availability of facilities for provision of signal functions

	Colombo								Gampaha								Kalutara					
	TH CSHW	TH DMH	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH RAGAMA	DGH GAMPAA	DGH NEGOMBO	BH WATHUPTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		GH KALUTARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	
At least one parenteral antibiotic	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
At least one parenteral uterotonic drug	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Magnesium sulphate	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Pethidine	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Adrenaline	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Vacuum Extractor	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Forceps	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Neonatal Ambu bag	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Neonatal sucker	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
At least one specialist obstetricians	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
At least one MO BTS	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
At least one MO trained in anaesthesia	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
At least two specialist obstetricians	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
At least three MO BTS	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
At least two or more persons trained in anas	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
House officers available	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Medical officers	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Nursing officers	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Midwives	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Electricity	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Continuous water	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Functional generator	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Functional operating theatre	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Functional blood bank	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Training issues																						
Manual removal of the placenta	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Assisted Vaginal Deliveries	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	
Neonatal Resuscitation	●	●	●	●	●	●	●		●	●	●	●	●	●	●		●	●	●	●	●	

● Available

● Not Available

● Not relevant

Table 3.8: Institutions that have potential ability to provide signal functions by district

	COLOMBO							GAMPAHA							KALUTARA					
	TH CSHW	TH DMH	TH Colombo South	BH Avissawella	BH Homagama	BH Mulleriyawa	DH Piliyandala	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	BH Mirigama	BH Kiribathgoda	DH Minuwangoda	GH Kalutara	BH Panadura	BH Horana	BH Pimbura	DH Mathugama	DH Beruwala
Administering of parenteral antibiotics																				
Administering of parenteral oxytocins																				
Administering of parenteral anti convulsants																				
Performing manual removal of placenta																				
Removal of retained products of conception																				
Performing assisted vaginal delivery																				
Performing neonatal resuscitation																				
Performing obstetric surgeries																				
Performing blood transfusions																				

Capable of performing the function
 Not capable of performing the function
 Not relevant

Table 3.8 shows that the Colombo district has 6 institutions capable of providing all 9 functions and be called a CEmONC facility. There are 4 such CEmONC facilities in Gampaha and 3 in Kalutara. In addition the province has 1, 2, 1 institution in Colombo, Gampaha and Kalutara districts respectively that are capable of the first 4 functions.

Emergency functions should be available for patients on a 24*7 basis. Based on criteria given in chapter 2, the ability of institutions to provide EmONC services on a 24*7 basis was examined. This is shown in table 3.9.

Table 3.9: Institutions that are able to provide CEmONC services on 24*7 by district

	COLOMBO							GAMPAHA							KALUTARA					
	TH CSHW	TH DMH	TH Colombo South	BH Avissawella	BH Homagama	BH Mulleriyawa	DH Piliyandala	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	BH Mirigama	BH Kiribathgoda	DH Minuwangoda	GH Kalutara	BH Panadura	BH Horana	BH Pimbura	DH Mathugama	DH Beruwala
Administering of parenteral antibiotics																				
Administering of parenteral oxytocins																				
Administering of parenteral anti convulsants																				
Performing manual removal of placenta																				
Removal of retained products of conception																				
Performing assisted vaginal delivery																				
Performing neonatal resuscitation																				
Performing obstetric surgeries																				
Performing blood transfusions																				

Capable of performing the function 24*7
 Not capable of performing the function 24*7
 Not relevant

Table 3.9 shows that the Colombo district has 5 institutions capable of providing 24*7 services, one institution that provides the first 7 functions and one providing the first 4 functions. Gampaha has 4 institutions that provide all 9 signal functions 24*7 and 2 which provide the first 4 functions. Kalutara has 2 institutions capable of providing CEmONC functions 24*7 and one which provides the BEmONC

functions and one institution that can provide the first 4 functions.

3.3 Availability of EmONC services (Indicator 1)

The first indicator examines the availability of EmONC services in terms of the population.

Table 3.10 (a) : Availability of EmONC facilities by district based on services provided during the 3 months prior to survey

	<i>Colombo</i>	<i>Gampaha</i>	<i>Kalutara</i>	<i>Western Province</i>
Population in 2012	2,323,826	2,298,588	1,214,880	5,837,294
Geographic extent in km ²	699	1387	1598	3684
Institutions which have provided all nine signal functions	4	4	3	11
Institutions estimated to have provided modified BEmONC services	0	1	1	2
Institutions estimated to have provided first 4 functions only	0	0	0	0
Availability of institutions that have provided modified BEmONC services per 500,000 population	0.0	0.2	0.4	0.17
Availability of institutions that have provided the first 4 signal functions only per 500,000 population	0.0	0.0	0.0	0.0
Availability of institutions that have provided all nine signal functions per 500,000 population	0.86	0.87	1.23	0.94
Area per modified B -EmONC facility (km ²)	NA	1387	1598	1842
Area per C-EmONC facility (km ²)	175	347	533	335
Area per EmONC facility (km ²) (CEmONC + Modified BEmONC)	175	277	400	283

Table 3.10 (b): Availability of EmONC facilities by district based on the potential ability of institutions for providing signal functions

	<i>Colombo</i>	<i>Gampaha</i>	<i>Kalutara</i>	<i>Western Province</i>
Population in 2012	2,323,826	2,298,588	1,214,880	5,837,294
Geographic extent in km ²	699	1387	1598	3684
Facilities which are able to provide modified BEmONC services	1	2	1	4
Facilities which are potentially able to provide C-EmONC services	6	4	3	13
Modified BEmONC facilities per 500,000 population	0.2	0.4	0.4	0.3
Institutions that have the potential ability to provide CEmONC services 500, 000 population	1.29	0.87	1.23	1.11
Institutions with modified BEmONC + CEmONC facilities per 500,000 population	1.49	1.27	1.63	1.41
Area per modified BEmONC facility (km ²)	699	694	1598	921
Area per CEmONC facility (km ²)	117	347	533	283
Area per EmONC facility (km ²) (CEm ONC + Modified BEmONC)	100	231	400	217

Table 3.10 (c): Availability of institutions that are able to provide CEmONC services on a 24*7 basis by district

	<i>Colombo</i>	<i>Gampaha</i>	<i>Kalutara</i>	<i>Western Province</i>
Population in 2012	2,323,826	2,298,588	1,214,880	5,837,294
Geographic extent in km ²	699	1387	1598	3684
Facilities which are able to provide 24*7 CEmONC services	5	4	2	11
Facilities which are able to provide 24*7 CEmONC if Blood bank services are upgraded	5	4	3	12
24*7 CEmONC facilities per 500,000 population	1.08	0.87	0.82	0.94
24*7 CEmONC facilities after improving blood bank services	1.08	0.87	1.23	1.03
Area per 24*7 CEmONC facility (km ²)	140	347	799	335
Area per 24*7 CEmONC facility after improving blood bank services (km ²)	140	347	533	307

The accepted minimum level for availability of EmONC services is; five EmONC facilities, for every 500,000 population, at least one of which should provide comprehensive care. The province does not have any BEmONC facilities. Districts of Colombo and Kalutara have sufficient number of CEmONC facilities per 500 000 population. Gampaha is unable to meet the required standard.

When 24*7 capability of CEmONC services is examined it is seen that only the Colombo district meets the standard and the province also falls short of meeting the requirement per 500 000 population.

3.4 Geographic distribution of EmONC facilities (Indicator 2)

Table 3.10b shows that the average area served by an institution capable of providing CEmONC services is smallest in Colombo followed by Gampaha and Kalutara. It is noted that in the Colombo district the facilities are crowded together. This indicator does not provide evidence about the spread of facilities. Average distances to services and travel times would be more meaningful and could be estimated using Geographic Information Systems.

3.5.1 Proportion of births in EmONC facilities (Indicator 3)

This is a very crude indicator of utilisation of EmONC services by pregnant women. The optimum long term goal must be to ensure that all women deliver in a place where emergency services are available. The indicator is calculated as the proportion of all births in an area that takes place in EmONC facilities. The denominator is the estimated live births in the area regardless of where the birth takes place. In the current analysis the indicator is calculated using RGs data.

Table 3.11, shows the proportion of deliveries occurring in CEmONC facilities range from 81% in Colombo to 96% in Kalutara. When births in government hospitals are used as the denominator it is seen that 98% of the deliveries in the Colombo district take place in CEmONC facilities. These proportions are reduced when the proportion of births occurring facilities capable of providing 24*7 services are considered.

Table 3.11: Proportion of births occurring in CEmONC facilities by district and province

	<i>Colombo</i>		<i>Gampaha</i>		<i>Kalutara</i>		<i>Western Province</i>	
	<i>No.</i>	<i>%</i>	<i>No.</i>	<i>%</i>	<i>No.</i>	<i>%</i>	<i>No.</i>	<i>%</i>
<i>Number of births registered by RG in 2011</i>	54,888		27,457		16,947		99,292	
<i>In CEmONC facilities</i>	44513	81.1	23934	87.2	16417	96.8	84,864	85.5
<i>In 24*7 CEmONC facilities</i>	44271	80.6	23934	87.2	12880	76.0	81085	81.7
<i>In 24*7 CEmONC facilities once blood bank services upgraded</i>	44271	80.6	23934	87.2	16417	96.8	84622	85.0

3.5 Utilization of EmONC facilities

Indicators 3 and 4 are on utilisation of the available EmONC facilities. They examine if the EmONC services serve a reasonable proportion of women and if the women who need the services i.e. those with complications receive the services.

This is a very crude indicator of utilisation of EmONC services by pregnant women. The optimum long term goal must be to ensure that all women deliver in a place where emergency services are available. There is no internationally advocated minimum requirement for this indicator. This has to be developed in consultation with stakeholders nationally

3.5.2 Met need for EmONC care (Indicator 4)

'Met need' is the proportion of women with major direct obstetric complications treated at EmONC facilities and is calculated as explained in table 2.2. It is estimated that about 15% of women develop a direct obstetric complication that may need emergency care. The direct obstetric complications considered in constructing this indicator are: haemorrhage (ante partum and post-partum), prolonged obstructed labour, postpartum sepsis, complications of abortions, severe pre-eclampsia and eclampsia, ectopic pregnancy and ruptured uterus.

The estimate based on the above assumption forms the denominator of this indicator. The numerator for this indicator in the present study is the actual number of women who were diagnosed as having a major direct obstetric complication in the EmONC facilities during the one month prospective survey of morbidity.

The minimum acceptable level of this indicator is 100% i.e. all women requiring EmONC care should receive such care. The table above (3.12) shows that nearly 75% of obstetric emergencies have received EmONC care in the Western Province.

3.6 Caesarean section (CS) as a proportion of all births (Indicator 5)

The proportion of caesarean sections may be considered as an indicator of utilisation of a lifesaving EmONC function. Although literature suggests that both very low and very high rates of caesarean sections may be unacceptable there is no empirical evidence on which to base an optimum proportion and this has provoked much debate. Pending further research based evidence the range of 5 – 15% is advocated by WHO. The threshold suggested by the RCOG is 25%.

In the present study data on caesarean sections are available from 4 sources. The proportion of sections by institution is available for 2010 from routine registers and for the quarter preceding data collection. Data was extracted from 10 CS using past records and information is also available from the prospective morbidity survey.

Table 3.13 shows the caesarean section rates for the province and the districts. These percentages are based on the registered births in each district.

Table 3.12: 'Met need' for EmONC care by district and province

	<i>Colombo</i>	<i>Gampaha</i>	<i>Kalutara</i>	<i>Western Province</i>
<i>Number of births registered by RG in 2011</i>	54,888	27,457	16,947	99,292
<i>Average number of registered births per month (RG)</i>	4222	2112	1304	7638
<i>Estimated number of direct obstetric complications</i>	633	317	196	1146
<i>Number of direct obstetric complications observed</i>	500	237	121	858
<i>'Met need' of EmONC (%)</i>	78.9	74.8	61.9	74.9

(Direct Obstetric Complications for Colombo district is without the data from Sri Jayawedhanapura TH)

Table 3.13: Caesarean section as proportion of births registered in each district and province

	<i>Colombo</i>	<i>Gampaha</i>	<i>Kalutara</i>	<i>Western Province</i>
	<i>No.</i>	<i>No.</i>	<i>No.</i>	<i>No.</i>
<i>Number of births per quarter for 2011 (based on RG data)</i>	13722	6864	4237	24823
<i>Total caesarean sections performed for quarter from birth register in 2011</i>	4123	1960	1428	7511
<i>Caesarean sections as proportion of all births – based on RG data</i>	30.0	28.6	33.7	30.3

(Data for Colombo district is without births from Sri Jayawedhanapura TH)

The caesarean section rate for Colombo is 30%, the rate for Gampaha and Kalutara being marginally lower and higher respectively. Calculations based on deliveries in government hospitals show a higher rate suggesting that the low rate may be because the RG data includes deliveries in the private sector but the numerator does not include the sections in that sector.

3.7 Measures of quality of services

Having examined the availability and utilisation of EmONC services the next three indicators examine some aspects of the quality of services. Measurement of quality of services is a complex process and the three indicators provide only a very crude measure of this aspect.

3.7.1 Direct obstetric case fatality rate (Indicator 6)

Case fatality rate is an indicator of the quality of care in addressing an obstetric emergency. The indicator

is defined as the proportion of women admitted to an EmONC facility with major direct obstetric complications, or who develop such complications after admission, and die before discharge. Direct causes of deaths are those 'resulting from obstetric complications of the pregnant state (pregnancy, labour, and puerperium), from interventions, omissions, incorrect treatment, or from a chain of events from any of the above'.

The last audited maternal death data were available for 2010. Therefore the number of women admitted to a facility, capable of providing EmONC services with a major direct obstetric complication needed to be estimated for the year 2010. This was carried out using the morbidity rates calculated for the different levels of institutions for the year 2011 based on the he prospective morbidity survey undertaken as part of the current study and the number of births in 2010 reported by the government institutions within the province capable of providing EmONC services.

Table 3.14: Direct obstetric case fatality rate by district and province

	<i>Colombo</i>	<i>Gampaha</i>	<i>Kalutara</i>	<i>Western Province</i>
	<i>No</i>	<i>No</i>	<i>No</i>	<i>No</i>
<i>Number of maternal deaths due to direct obstetric complications in 2010</i>	3	1	3	7
<i>Estimated number of direct obstetric complications for 2010</i>	6559	3397	1506	11462
<i>Direct obstetric case fatality rate for 2010 (%)</i>	0.046	0.029	0.199	0.061

The maximum acceptable level recommended for this indicator is a case fatality rate less than 1%. Table 3.14 shows that the case fatality rate for direct obstetric causes is well below the maximum acceptable level in both districts and the province as a whole. The case fatality rate is seen to be higher in the Kalutara district but without information on the condition of women on arrival at the facilities it cannot be inferred that the higher rate indicates poor quality of care.

3.7.2 Intrapartum and very early neonatal death rate (Indicator 7)

This indicator sheds light on the quality of intrapartum care for foetuses and new-borns delivered in EmONC facilities. Early neonatal deaths are defined as neonates born at term, who could not be resuscitated or for whom the resuscitation was not available or who had a specific birth trauma and died within the first 24 hours of life. In the present study information on early neonatal deaths are not available. Therefore the intra-partum death rate is calculated. The numerator is the reported number of still births in EmONC facilities and the denominator is the number of births for the quarter in the same institutions expressed as per 1000 births. Data for the same quarter in 2011 are used.

The total stillbirth rate is lowest in Gampaha and highest in Kalutara. In Colombo the percentage of macerated still births are higher than the fresh still births while the reverse is seen in Gampaha and Kalutara.

3.7.3 Proportion of deaths due to indirect causes in EmONC facilities (Indicator 8)

With increasing proportions of maternal deaths being attributed to indirect causes of death this is an important indicator. This reflects the shared medical care available for co-morbidities during pregnancy i.e. medical services other than EmONC services that are necessary to make pregnancy safer.

In calculating this indicator, all maternal deaths due to indirect causes in EmONC facilities within a specified period was taken as the numerator and all maternal deaths in EmONC facilities during the same period was taken as the denominator. Indirect causes of death results from 'previous existing diseases or diseases that developed during pregnancy and which was not due to direct obstetric causes, but which was aggravated by the physiological effects of pregnancy'.

Table 3.15: Stillbirth rate per 1000 births by district and province

	Colombo		Gampaha		Kalutara		Western Province	
	No	rate	No	rate	No	rate	No	rate
<i>No of births</i>	10058		6130		3898		20086	
<i>Fresh still births ≥ 2.5 Kg</i>	12	1.2	8	1.3	4	1.0	24	1.2
<i>Fresh still births < 2.5 Kg</i>	10	1.0	9	1.5	3	0.8	22	1.1
<i>Total fresh still births</i>	22	2.2	17	2.8	7	1.8	46	2.3
<i>Macerated still births</i>	32	3.2	13	2.1	0	0.0	45	2.2
<i>Unspecified</i>	0	0.0	0	0.0	21	5.4	21	1.0
<i>Total still births</i>	54	5.4	30	4.9	28	7.2	112	5.6

Table 3.16: Proportion of deaths due to indirect causes in CEmONC facilities

	<i>Colombo</i>	<i>Gampaha</i>	<i>Kalutara</i>	<i>Western Province</i>
<i>Number of maternal deaths due to indirect causes at EmONC facilities in 2010</i>	1	7	4	12
<i>Total number maternal deaths at EmONC facilities in 2010</i>	5	9	9	23
<i>Proportion of deaths due to indirect causes in 2010</i>	20.0	77.8	44.4	52.2

Gampaha (0.87) and in Kalutara it is more than half (0.57). There is no accepted level identified for this indicator. Because of the small numbers it is more meaningful to examine these at national level.

3.8. Summary of EmONC indicators 1-8

Table 3.17 and figure 3.1 summarises the EmONC indicators for the Province.

Figure 3.1 is a summary matrix based on the table above in a visual format which compares the performance of the different districts on a given indicator. The colour codes in each square indicate the level of accomplishment and are divided as given below in descending order of shades.

- Dark green - achievement of the intended goal,
- Next shade - below the intended goal but above the national average
- Next lighter shade - performance below the intended goal and below the national average, but better than the lower benchmark.
- Lightest shade lower than the lower benchmark

The values for each indicator are given in the key below the table. Where there were no standard benchmarks which could be used set nationally or internationally, cut off values were based on expert consensus.

Table 3.17: Summary of EmONC indicators 1-8 by district

District	Availability of services			Accessibility of services			Utilisation of services			Quality of care				
	Indicator 1			Indicator 2			Indicator 3			Indicator 4	Indicator 5	Indicator 6	Indicator 7	Indicator 8
	C_EmONC capable / 500000 population	24*7 C_EmONC capable/ 500000 population	24*7 C_EmONC capable/ 500000 population once BB services upgraded	Area C_EmONC capable facility km ²	Area 24*7 C_EmONC capable facility km ²	Area 24*7 C_EmONC capable facility once BB services upgraded km ²	% Deliveries in CEmONC capable facilities	% Deliveries in 24*7 C_EmONC capable facilities	% Deliveries in 24*7 C_EmONC capable facilities once BB services upgraded	Met need of EmONC care	Caesarean Section Rate	Direct Obstetric Case Fatality Rate	Total still birth rate	Proportion of deaths due to indirect causes - 2010
Colombo	1.29	1.08	1.08	117	140	140	81	81	81	79.0	30	0.046	5.4	20.0
Gampaha	0.87	0.87	0.87	347	347	347	87	87	87	74.8	28.6	0.029	4.9	77.8
Kalutara	1.23	0.82	1.23	533	799	533	97	76	97	61.7	33.7	0.199	7.2	44.4
Sri Lanka	1.65	0.79	0.96	979	2050	1682	88	62	73	80.0	27.4	0.151	6	31.6

The cut offs for CEmONC facilities per 500,000 population and the 24*7 CEmONC facilities were decided upon based on the fact that the country as a whole does not meet the criteria for 5 EmONC facilities per 500,000 population and that the people's preference appear to be to deliver in a specialist unit that is capable of comprehensive EmONC services. As such the criterion of one CEmONC facility per 500,000 population is considered inappropriate. The current national average is 1.65 CEmONC facilities per 500,000 populations (range 5.42-0.87 per 500,000 population). Considering the current birth rate and the fact that 5 EmONC facilities are advocated internationally for adequate availability, it was decided that 3 CEmONC facilities per 500,000 population should be considered as the immediate goal. Considering the current low availability of 24*7 CEmONC facilities it was decided that there should be at least one institution per 500,000 populations that is capable of 24*7 CEmONC services and should be a goal to work towards in the immediate future.

3.9. New indicators developed in the current study

The analysis presented above shows that indicators one and two has to be examined in the Sri Lankan context. It is also seen that the community preference to deliver in a specialist unit is high and the proportion delivering in such units have been increasing over time. As such aiming to provide 4 BEmONC or modified B-EmNOC facilities per 500 000 population would be of limited value in planning future services. Therefore the current analysis refined indicator two and identified some new indicators to help in planning services for the future. These are presented in sections 3.9-3.12.

3.9.1 Refining indicator 2 (Geographic distribution of EmONC facilities)

Indicator two (as defined in section 3.2) calculated as the average land area covered per facility per district is a crude indicator of access to services and does not identify spatial clustering of institutions.

Figure 3.1: Equity Assessment Matrix of the 8 EmONC indicators


District	C_EmONC capable / 500000 population	24*7 C_EmONC capable/ 500000 population	24*7 C_EmONC capable/ 500000 population once BB services upgraded	Area C_EmONC capable facility km ²	Area 24*7 C_EmONC capable facility km ²	Area 24*7 C_EmONC capable facility once BB services upgraded km ²	% Deliveries in CEmONC capable facilities	% Deliveries in 24*7 C_EmONC capable facilities	% Deliveries in 24*7 C_EmONC capable facilities once BB services upgraded	Met need of EmONC care	Caesarean Section Rate	Direct Obstetric Case Fatality Rate	Total still birth rate	Proportion of deaths due to indirect causes - 2010
Colombo	1.29	1.08	1.08	117	140	140	81	81	81	79.0	30.0	0.046	5.4	20.0
Gampaha	0.87	0.87	0.87	347	347	347	87	87	87	74.8	28.6	0.029	4.9	77.8
Kalutara	1.23	0.82	1.23	533	799	533	97	76	97	61.7	33.7	0.199	7.2	44.4
Sri Lanka	1.65	0.79	0.96	979	2050	1682	88	62	73	80.0	27.7	0.151	6.0	31.6
Standard														
	>=3	>=1	>=1	< 500	< 500	< 500	> 75	> 75	> 75	> = 100	5 - 15	< 0.151	< 6	< 25
	3.0 - 1.65	0.8 - 0.9	0.8 - 0.9	501- 1500	501- 1500	501- 1500	74 - 50	74 - 50	74 - 50	99 - 75	15 - 25	.151 - 0.5		25.1 - 50.0
	1.64 - 1.00	< 0.8	< 0.8	1501 - 2500	1501 - 2500	1501 - 2500	49 - 25	49 - 25	49 - 25	74 - 50	> 25.1	0.51- 1.0		50.1 - 75.0
	< 0.99	0	0	>250 1	>250 1	>250 1	< 25	< 25	< 25	< 50	< 5.00	> 1.00	> 6	> 75.1

3.9.1 (a): Percentage area falling outside a defined buffer zone round CEmONC facilities

To further refine indicator 2 the current study examined the area falling outside a defined buffer zone round a CEmONC facility. The figure of 30 km was decided upon as an average distance that could be travelled in 45 minutes to one hour to reach a CEmONC facility using common modes of transport available to a majority of the population.

It must be noted that the actual time from emergency to receiving appropriate services would be more than 45 minutes taking in to consideration the time lags between making the decision to seek care and starting travel as well as time between arrival in an institution and receiving care.

Figure 3.2 shows facilities capable of providing CEmONC services and those with 24*7 cover with buffers of 30 km. radius and the area that is not covered by such services.



The percentage area falling outside the 30 km buffer zone shown in figures 3.2 are given in table 3.18.

Table 3.18: Area per district falling outside a 30 km. buffer zone from CEmONC and 24*7 CEmONC facilities

District	Land area without forest cover Km ²	Based on institutions that are potentially able to provide CEmONC services			Based on institutions able to provide 24*7 CEmONC services			Based on institutions able to provide 24*7 CEmONC services (after correction of BTS)		
		Sq. Km outside buffer	% land area outside buffer *	% from total unserved area for the country	Sq. Km outside buffer	% land area outside buffer	% from total unserved area for the country	Sq. Km outside buffer	% land area outside buffer	% from total unserved area for the country
Colombo	679.7	0	0	0	0	0	0	0.0	0.0	0.0
Gampaha	1386	0	0	0	0	0	0	0.0	0.0	0.0
Kalutara	1645.4	37.1	2.3	0.5	37.1	2.3	0.1	37.1	2.3	0.2
Sri Lanka	59237.7	7991.6	13.5	100	27044.7	45.7	100	20085	30.5	100.0

*In this calculation the forest areas have been left out however, the data on area under agriculture were not available.

The tables show that the area falling outside a 30km buffer zone round facilities that have the potential ability to provide CEmONC services is 13.5% at the national level and only 2.3% of land area of Kalutara district lie outside such facility. This proportion is same when facilities able to provide CEmONC services on a 24*7 basis are examined.

3.9.1 (b): Percentage population falling outside a defined buffer zone round CEmONC facilities

In interpreting the above indicator it must be noted that there is considerable overlap between areas served by institutions and spill over in to neighbouring districts. Therefore in the current study this was further refined by overlaying the population figures from Census 2012, at Divisional Secretariat level and calculating the population without EmONC cover i.e. population outside the 30 kilometre buffer zone and this is presented in table 3.19.

Table 3.19: Population without access to CEmONC and 24*7 CEmONC facility cover using a 30 km buffer zone.

District	Population 2012	Based on institutions that have the potential ability to provide CEmONC services			Based on institutions able to provide 24*7 CEmONC services			Based on institutions able to provide 24*7 CEmONC services after correction of BTS		
		Number not having access to service	% not having access	% from total without access	Number not having access to service	% not having access to service	% from total without service	Number not having access to service	% not having access to service	% from total without access
Colombo	2309809	0	0	0	0	0	0.0	0	0.0	0.0
Gampaha	2294641	0	0	0	0	0	0.0	0	0.0	0.0
Kalutara	1217260	8026	0.7	1.4	8026	0.7	0.2	8013	0.7	0.4
Sri Lanka	20,263,723	582387	2.9	100.0	3913803	19.3	100.0	2065355	10.2	100.0

Since the population density varies between districts a more sensitive indicator for planning purposes may be the percentage population left out of reach of services. It is seen that only 3% of the total population in the country lie outside a 30 km buffer zone from a facility capable of CEmONC services while almost all the population of Western Province live within the 30km buffer zone for such facility.

This information could be further refined by calculating the proportion of the population falling outside a predetermined buffer based on average distances to services from human habitations and travel times to reach a facility. However, updated spatial data necessary for such a calculation were not available for the whole country at the time of analysis.

3.10 Availability and accessibility

Availability and accessibility have to be examined together for meaningful interpretation and planning. Figures 3.3 and 3.4 shows the relative position of the district in relation to other districts.

A radius of 30 km corresponds to a square area of 3000 km² approximately. It is seen from the diagram that where institutions capable of providing comprehensive emergency services are concerned there is no district where availability is below 1 facility per 500 000 population and the area served is more than 3000 km².

Figure 3.3: Scatter plot of CEmONC facilities by indicators one and two




Figure 3.4: Scatter plot of 24*7 CEmONC facilities by indicators one and two




Figure 3.4 highlights that 24*7 CEmONC services are deficient by both indicators in a number of districts. The figures clearly show the necessity to develop a composite indicator which would examine both accessibility and availability together.

The study developed the following indicators as a composite of indicator one and two to address such issues.

3.11 Composite indices developed in the study

Three composite indices which take into account availability and accessibility were developed in the study to help with prioritisation of districts for planning purposes and are presented in this section.

3.11.1 Accessibility Index for a Specialized hospital – (AISH)

The accessibility index for a specialized hospital is a composite index between representing the ratio between the average number of deliveries per specialized hospital (numerator) and the population density of a district (denominator).

AISH = Average number of deliveries per specialized hospital / population density of a district

In the spectrum of AISH, lower values represent higher accessibility and higher values represent lower accessibility.

3.11.2 Accessibility Index for a specialized unit – (AISU)

The accessibility index for a specialized unit is a composite index representing the ratio between the average number of deliveries per specialized unit (numerator) and the population density of a district (denominator).

AISU = Average number of deliveries per specialized unit / population density of a district

In the spectrum of AISU, lower values represent higher accessibility and higher values represent lower accessibility.

3.11.3. Accessibility index for a Specialist Obstetrician (AISO)

The accessibility index for a specialist obstetrician is a composite index between representing the ratio between the average number of deliveries per

specialist obstetrician (numerator) and the population density of a district (denominator).

AISO = Average number of deliveries per specialist obstetrician / population density of a district

In the spectrum of AISO, lower values represent higher accessibility and higher values represent lower accessibility.

Table 3.20 highlights the fact that the number of units available in the country (117) is sufficient to meet the norm (105) but there is inequity in distribution. The average number of deliveries per specialist obstetrician is highest in the Kalutara district and it is more than the national average.

Figure 3.5 summarises some of the indicators important for planning services.

3.12. Ministry of Health norms by district

In addition to the above, the current analysis examined the availability of specialists in the districts per population and area. This is given in Table 3.20. The Ministry of Health norm is to have 300 deliveries per unit per month.

Table 3.20: Specialist obstetricians per district according to Ministry of Health norms

Province	District	Population 2012	Total Births in the district in 2011 (RG)	No. of facilities with specialist care	No of specialist units in the district	No of specialist units required to meet the norm	No of Specialist Obstetricians available	Specialist units per 500000 population	Specialist obstetricians per 500000 population	Average number of deliveries per Specialist Obstetrician per month
Western province		5837294	99292	13	35	28	43	3.00	3.68	192
	Colombo	2323826	54888	6	20	15	25	4.30	5.38	183
	Gampaha	2298588	27457	4	9	8	12	1.96	2.61	191
	Kalutara	1214880	16947	3	6	5	6	2.47	2.47	235
Sri Lanka		20 277 597	343 384	69	117	105	138	2.59	3.40	207

Figure 3.5: Summary of indicators 1 and 2, new indicators and utilisation indicators by district

District	WHO Indicators 1 & 2				New Indicators						Utilization indicators				
	CEmONC / 500000 population	24*7 CEmONC / 500000 population	Area CEmONC facility	Area 24*7 CEmONC facility	Accessibility index for SH	Accessibility index for a SU	Accessibility index for a Specialist Obstetrician	% population outside CEmONC	% population outside 24*7 CEmONC	% population outside 24*7 CEmONC once BB services upgraded	Specialist unit / 500000 population	Specialist Obstetrician / 500 000 population	Average deliveries / Specialist VOG /month	Average deliveries / Specialist hospital /month	Average deliveries / Specialist unit /month
Colombo	1.29	1.08	117	140	2.75	0.83	0.66	0	0.0	0.0	4.3	5.38	183	762	229
Gampaha	0.87	0.87	347	347	4.41	1.84	1.38	0	0.0	0.0	1.96	2.61	191	572	254
Kalutara	1.23	0.82	533	799	7.43	3.72	3.72	0.7	0.2	0.4	2.47	2.47	235	471	235
Sri Lanka	1.65	0.79	979	2050	16.11	9.50	8.52	2.9	19.3	10.2	2.59	3.40	207	415	245
Green	>=3	>=1	< 500	< 1000	< 5	< 5	< 5	0.0	0.0	0.0	>= 3	> 3	=< 250	=< 500	=< 250
Yellow	3.0 - 1.65	.8 - .9	501- 1000	< 2050	5.1- 10.0	5.1 - 10	5.1 - 10	0.1 - 5.0	0.1 - 5.0	0.1 - 5.0					
Orange	1.64 - 1.00	< 0.8	1001- 1500	> 2050	10.1- 20.0	10.1 - 20.0	10.1 - 20.0	5.1 - 10.0	5.1 - 10.0	5.1 - 10.0					
Red	< 0.99	0	>1501	0	> 20.1	> 20.1	> 20.1	> 10.1	> 10.1	> 10.1	< 3	< 3	>= 251	>= 501	>= 251

In deciding the cut off value for average number of deliveries per specialist obstetrician the figure of 250 was decided upon taking the national average of 207 in to consideration.

In deciding on a cut off for the average deliveries per specialist hospital 500 (250*2) was decided upon since a minimum of two specialists should be available for the provision of 24*7 CEmONC care.

Although many of the TH/PGH type of institution may have more than two specialists, the total number of institutions in this group is small.

The above figure and the numbers there in are presented so that they can be used to initiate a discussion and national consensus on standards for CEmONC care.

Chapter 4

Selected Care Practices

4.1 Caesarean sections

Data on caesarean sections were available from multiple sources:

- H830 for 2010 and the quarter preceding data collection 2011,
- Ten Bed Head Tickets of patients who have had Caesarean sections,
- Prospective morbidity survey (4 weeks).

4.1.1 Institutional caesarean section rates

In table 4.1 the caesarean section rates are calculated based on the actual number of deliveries that took place in the institutions. Hospital rates are difficult to interpret, since they depend on the case mix seen in the institution as well as preferences of providers and clients.

Table 4.1: Caesarean section, rates by institution district and province

Hospital	COLOMBO						GAMPAHA				KALUTARA			Colombo District	Gampaha District	Kalutara District	Western Province
	TH Castle Street	TH De Soysa	TH Colombo South	BH Awissawella	BH Homagama	BH Mulleriyawa	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	DGH Kaluthara	BH Panadura	BH Horana				
<u>Based on H830 return for 2010</u>																	
Total reported deliveries/year	16445	8935	8663	4579	2551	70	8813	5236	3290	3350	8014	3913	1230	41243	20689	13157	75089
Total caesarian sections/year	5765	2461	2741	1676	732	11	3018	1672	1153	1243	3050	1431	605	13386	7086	5086	25558
Institutional section rate	35.1%	27.5%	31.6%	36.6%	28.7%	15.7%	34.2%	31.9%	35.0%	37.1%	38.1%	36.6%	49.2%	32.5%	34.3%	38.7%	34.0%
<u>From the birth register for a quarter of 2011</u>																	
Total reported deliveries/quarter	3872	2272	1993	1226	695	70	2123	1391	1408	1208	2177	1041	884	10128	6130	4102	20360
Total caesarian sections/quarter	1304	365	633	494	199	11	740	411	477	332	762	357	332	3006	1960	1451	6417
Institutional section rate	33.7%	16.1%	31.8%	40.3%	28.6%	15.7%	34.9%	29.5%	33.9%	27.5%	35.0%	34.3%	37.5%	29.7%	32.0%	35.4%	31.5%

(* District and provincial rates calculated based on deliveries at specialized units only)

The table above (4.1) shows that the section rates in the two major hospitals for women in the country and all other institutions in the Colombo district except one has declined or remained the same from 2010 to 2011. Similar decline is seen in Gampaha and Horana districts too.

The morbidity survey provided information on the proportion of caesarean sections by category. It is noted that the majority are emergency sections except in one institution where the proportions appear to be reversed.

Table 4.2 shows that in the 3 teaching hospitals in the Colombo district the percentage of emergency sections is greater than the proportion of elective sections. In the two base hospitals in the district the pattern is reversed.

In the teaching hospital in Gampaha the proportion of elective sections is less than the proportion of emergency sections. The pattern is reversed in the two DGHs and the BH.

In Kalutara the DGH has a lower proportion of elective sections while in the two base hospitals the pattern is reversed.

4.1.2 Indications for sections

Details of caesarean sections obtained from studying 10 BHTs are presented in tables 4.3 and 4.4.

Table 4.3 lists the indications for sections, which were obtained from the 10 BHTs perused per unit and the analysis is presented for the province.

A previous caesarean section is the indication for section in 58% of women.

Table 4.2: Category of caesarean sections by institution and by districts based on one month prospective survey

	WESTERN PROVINCE													
	GAMPAHA					KALUTARA				COLOMBO				
	TH COLOMBO NORTH	DGH GAMPAHA	DGH NEGOMBO	BH WATUPITIWALA		DGH KALUTARA	BH PANADURA	BH HORANA		TH CASTLE STREET	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA
Caesarean Section - Elective	67	98	75	75		80	67	33		94	82	62	87	35
Caesarean Section - Emergency	70	55	54	62		95	54	21		112	89	127	73	30
Total Caesarean Section	137	153	129	137		175	121	54		206	171	189	160	65
Total deliveries	546	497	436	465		601	341	206		663	434	541	362	188
CS as a percentage of total deliveries	25.1	30.8	29.6	29.5		29.1	35.5	26.2		31.1	39.4	34.9	44.2	34.6
Percentage of elective sections	48.9	64.1	58.1	54.7		45.7	55.4	61.1		45.6	48.0	32.8	54.4	53.8
Percentage of emergency sections	51.1	35.9	41.9	45.3		54.3	44.6	38.9		54.4	52.0	67.2	45.6	46.2

Table 4.3: Indications for Caesarean Section

	Number	Percentage
No of CS reviewed	170	
Previous caesarian section	98	58%
Fetal distress	82	48%
Failed induction	39	23%
Any other obstetric complication	28	16%
Pre-eclampsia/ Eclampsia	22	13%
Cephalopelvic disproportion/dystocia/ failure to progress/ failed vacuum extraction or forceps	19	11%
Any other maternal medical complication	17	10%
Breech	11	6%
Other mal presentations	11	6%
Suspected fetal growth impairment	10	6%
Gestational age 41 completed weeks or more	9	5%
Multiple pregnancy	9	5%
Any other fetal indication	7	4%
Elderly mother	6	4%
Subfertility	5	3%
Elderly primi	4	2%
Maternal request	3	2%
Large baby	3	2%
Previously repaired vesico-vaginal or recto-vaginal fistula	2	1%
Tubal ligation/sterilization	1	1%
Previous uterine surgery	1	1%

The data were further analysed to examine the use of partographs among women who had emergency caesarean sections. A partograph had been

maintained only for 90.2%, 60% and 50% of the emergency sections in Colombo, Gampaha and Kalutara district respectively.

4.1.3 Selected care practices for caesarean section

Table 4.4: Service provision for caesarean sections

	COLOMBO					GAMPHA				KALUTARA			Colombo District	Gampaha District	Kalutara District	Western Province
	TH Castle Street	TH De Soysa	TH Colombo South	BH Awissawella	BH Homagama	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	DGH Kaluthara	BH Panadura	BH Horana				
<i>C/S by selected characteristics</i>																
No. of CS reviewed	50	46	30	20	20	30	20	20	20	20	20	20	166	90	60	316
Percentage decided by a specialist obstetrician	96	88.2	100	100	95	100	100	85	100	95	100	100	94	96.7	98.3	95.5
Percentage performed by a specialist obstetrician	10	6.5	6.7	0	20	0	50	0	0	0	15	0	8.4	11.1	5	8.5
Percentage performed by non-specialist MO	90	91.3	93.3	100	80	100	50	100	100	100	85	100	91	88.9	95	91.2
Proportion of CS transferred from other institutions	4		6.7	0	0	3.3	10	0	0	0	0	0	2.4	3	0	1.9
Percentage CS under general anaesthesia	0	0	3.3	0	5	0	5	10	0	0	5	5	1.2	2.2	3.3	1.9
Percentage CS under other anaesthesia (spinal and epidural)	100	95.7	96.6	100	95	100	95	90	100	100	95	95	97.6	96.7	96.7	97.1
percentage of CS that received antacid prophylaxis	40	84.8	60	65	80	100	100	70	20	85	100	100	63.9	75.6	48	70.3
Percentage of CS administered prophylaxis antibiotics	66	80.4	93.3	95	50	100	100	100	100	15	10	0	76.5	100	57	80.4
Percentage CS with surgical complications	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Percentage with anaesthetic complications	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Mean stay at hospital after CS (Days)	5.1	3.8	4.2	4.5	4.9	3.4	3.1	3.4	3.8	2.6	3.2	4.1	4.5	3.8	3.3	3.9
Modal stay at the hospital after CS (Days)	4	4	3	5	3	3	3	3	4	2	3	3	3	3	3	3

In the majority (88%-100%) the decision for caesarean section has been taken by the specialist obstetrician. The proportion of section has been performed by the specialist obstetrician varies from 0-67%. Antacid prophylaxis varies from 20%-100% while use of prophylactic antibiotic use varies from 10% to 100%. The median hospital varies from 3-5 days, in the majority of the institutions the median is 3 days.

The data were further analysed to examine the use of partographs among women who had emergency caesarean sections. A partograph had been maintained for 70.6% of the emergency sections in the province while 90.2% in the Colombo district.

4.2. Use of partographs

The partograph use provides some insights in to monitoring of labour and may be considered as being indicative of quality of intra natal care. Table 4.5 gives details on selected aspects of partograph use in the three districts.

A majority of institutions in the province use the partograph. Non use is mainly in the smaller institutions where its use is even more important. The quality of recording is best in the Colombo district and recordings relatively poor in the Kalutara district.

Table 4.5 Use of the partograph by district

	Colombo								Gampaha						Kalutara					
	TH Castle Street	TH De Soyza	TH Colombo South	BH Awissawella	BH Homagama	BH Mulleriyawa	DH Piliyandala	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	BH Meerigama	BH Kiribathgoda	DH Minuwangoda	DGH Kaluthara	BH Panadura	BH Horana	BH Pimbura	DH Mathugama	DH Beruwala
Institutions using partographs																				
Number of partographs reviewed	25	24	10	10	5	0	0	15	10	5	10	0	0	0	10	5	5	0	1	0
	No.				%				No.				%		No.				%	
Partograph was used when in labour	79				94.0				39				97.5		20				95.2	
Charting started when admitted to LR	67				79.8				35				87.5		19				90.5	
Temperature recorded at least once	52				61.9				20				50.0		7				33.3	
Blood pressure recorded at least once	51				60.7				25				62.5		10				47.6	
Maternal pulse recorded at least once	81				96.4				26				65.0		10				47.6	
Foetal heart sounds recorded at least once	82				97.6				36				90.0		21				100.0	
Contractions recorded at least once	68				81.0				22				55.0		11				52.4	
Vaginal Ex. Findings recorded at least once	84				100.0				38				95.0		10				47.6	
Descent of the head recorded at least once	63				75.0				18				45.0		9				42.8	
State of membranes mentioned	70				83.3				29				72.5		8				38.1	
Alert line drawn	62				73.8				26				65.0		7				33.3	
Time of delivery filed in	83				98.8				25				62.5		11				52.4	
Findings marked elsewhere in the BHT	40				47.6				18				45.0		13				61.9	



Using a Partograph



Not using a Partograph

4.3 Management of labour

4.3.1 Induction of labour

Table 4.6: Induction of labour in specialist and non-specialist institutions in Western province

	Gampaha-Specialized		Gampaha-non Specialized		Kalutara – Specialized		Kalutara – non specialized		Colombo – Specialized		Colombo – Non Special		Western-Specialized		Western – Non Specialized	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Total BHT's Examined	2699		NI		1565		NI		3325		3		7589		3	
Total BHT's reviewed in Obstetric wards	2408	89.2	NI	NI	1492	95.3	NI	NI	2408	89.2	NI	NI	1492	95.3	NI	NI
-																
Documented details of labour	1514	76.3	NI	NI	712	61.7	NI	NI	1592	68.7	0	0.0	3818	70.0	0	0.0
Labor - spontaneous	788	52.0	NI	NI	269	37.8	NI	NI	814	51.1	0	0.0	1871	49.0	0	0.0
Labor - induced	726	48.0	NI	NI	443	62.2	NI	NI	778	48.9	0	0.0	1947	51.0	0	0.0
If induced																
Syntocinon	560	77.1	NI	NI	375	84.7	NI	NI	551	70.8	0	0.0	1486	76.3	0	0.0
prostaglandin (PG)	47	6.5	NI	NI	2	0.5	NI	NI	58	7.5	0	0.0	107	5.5	0	0.0
Synto + PG	60	8.3	NI	NI	34	7.7	NI	NI	92	11.8	0	0.0	186	9.6	0	0.0
Other and non-specified	59	8.1	NI	NI	32	7.2	NI	NI	77	9.9	0	0.0	168	8.6	0	0.0

Table 4.6 shows that in the province as a whole labour is induced in more than half the patients in specialised

units. The proportion is particularly high in the Kalutara district.

4.3.2 Selected practices related to labour

Table 4.7: Practices related to labour

	COLOMBO																GAMPAHA										KALUTARA								
	TH CSHW	TH CSHW	TH CSHW	TH DMH	TH DMH	TH DMH	TH DMH	TH Colombo South	TH Colombo South	TH Colombo South	BH AVISSAWELLA	BH HOMAGAMA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA	TH COLOMBO NORTH	TH COLOMBO NORTH	TH COLOMBO NORTH	DGH GAMPAHA	DGH GAMPAHA	DGH NEGOMBO	DGH NEGOMBO	BH WATHUPITIWALA	BH WATHUPITIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA		
Episiotomy																																			
The current policy of the unit regarding application of episiotomy																																			
Perform Episiotomy on all primies	✓	✓	✓	✓	✓	✓	✓	✓			✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓						✓	✓	✓	✓			
Perform episiotomy on all multies	✓							✓	✓	✓															✓										
Perform episiotomy on selected primies																																			
Perform episiotomy on selected multies		✓		✓	✓	✓	✓							✓			✓	✓	✓	✓	✓	✓	✓	✓					✓	✓	✓	✓	✓		
Local anesthesia to perform epis		✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✗	✗	✓	✓	✗	✗	✗	✗	✗	✗			✗	✓	✗	✗	✗	✗	✗	
Local anesthesia to suture	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✗		✓		✓	✓	✓	✓	✓	✓	
The person who performs the episiotomy																																			
HO/MO	✓	✓	✓		✓												✓	✓				✓	✓											✓	
Nursing officer	✓	✓	✓	✓	✓	✓		✓	✓			✓	✓	✓	✓		✓	✓				✓	✓	✓	✓		✓		✓	✓	✓	✓	✓		
Midwife					✓	✓																												✓	
The person who suture the episiotomy																																			
HO/MO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓			✓		✓	✓	✓	✓	✓	
Nursing officer																																			
Midwife																									✓										
Pain relief																																			
The pain relief policy for normal vaginal delivery																																			
Epidural		✓		✓	✓	✓																	✓							✓					
Pethidine	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓	✓				✓	✓	✓	✓	✓			✓	✓	✓	✓	✓	✓	
None															✓													✓						✓	✓
Active management of thirdstage of labour																																			
A clear policy on active management of third stage of labour is available	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓		✓	✓	✓	✗	✗	✗	
A written document giving policy / flow chart/ is displayed	✓	✓	✗			✓	✓	✓	✓	✓	✓	✗		✓	✗		✗	✓	✗	✗	✗	✗	✗	✗	✗	✓		✗	DNA	DNA	DNA	✗	✗	✗	
Oxytocin is given immediately after the delivery																																			
To all the cases	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓							✓	✓	✓		✓		✓	✓	✓	✓	✓	✓	
To selected cases															✓		✓	✓															✓	✓	

4.3.3 Referrals and average stay

Table 4.8:Referral practices and duration of stay by institution and district

	Colombo																					Gampaha												Kaluthara					
	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH DMH	TH DMH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH RAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		DGH KALUTHARA	DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA		
General referral																																							
plans/referrals/protocols for referrals displayed in the LR	✓	✓	✓	✓	✓	✗	✓	✗	✓		✓	✗	✗	✗	✓	✓	✓	✗	✗		✓	✗	✗	✗	✗	✓	✗	✗		✗		✓	✓	✓	✓	✗	✓	✗	
practice of ensuring service availability before transfer	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
Practice of informing the station before transfer	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	
practice of sending an appropriate staff member	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	
Duration of stay																																							
Duration of stay following ND - without an episiotomy (hrs)	12	24	24	24	24	24	24	24	24	24		24	24	24	24	24	24	24	24		24	24	24	24	24	24	24			24		24	24	24	24	24	24	24	
Duration of stay following ND - with an episiotomy (days)	1	1	1	1	1	1	1	1	1	1	2	1	1	1	1	1	1	1	1		1	1	1	1	1	1	1			2		1	1	1	1	1	1	1	
Duration of stay following assisted vaginal delivery (Days)	2	2	2	2	2	2	2	2	2	2	2	2	1	1	2	2	1	1	DNA		1	2	2	2	0	2	2	DNA	DNA		1	1	1	1					
Duration following a CS (Days)	2	3	3	4	4	3	3	3	2	3	4	3	2	4	3	3	3			2	3	3	4	3	3	4					3	2	3	3					

4.4 Availability of guidelines and protocols for EmONC

A list of ten guidelines and protocols relevant to EmONC services was agreed upon and information was collected on the availability of these in the wards and labour rooms.

Table 4.9 shows that the availability of protocols and guidelines is poor in the province, the worst affected being the Kaluthara district. The most widely available are the labour room guide, Pregnancy, childbirth, postpartum and newborn care guide, breast feeding booklet and the breast feeding flash cards.

Table 4.9: Availability of guide lines and protocols by institution

	COLOMBO														GAMPAHA										KALUTARA											
	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH DMH	TH DMH	TH DMH	TH DMH	TH DMH	TH COLOMBO SOUTH	TH COLOMBO SOUTH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA	TH COLOMBO NORTH	TH COLOMBO NORTH	TH COLOMBO NORTH	DGH GAMPAHA	DGH GAMPAHA	DGH NEGOMBO	DGH NEGOMBO	BH WATHUPITIYAWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATUGAMA	DH BERUWALA	
HSDP National Guidelines	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
Labour room management guide	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
Pregnancy, Child birth, Postpartum and Newborn Care Guide	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
Guidelines for the Management of HIV infection in Pregnancy in Sri Lanka	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
National guidelines for the prevention & management of malaria in pregnancy	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
Guidelines on immunization against Tetanus	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
Management of tuberculosis during pregnancy	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
Neonatal advanced life support protocol (wall chart)	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
Breast feeding booklet	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available
Breast feeding flash cards	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available	Available

Available

Not available

No Information

■ Available
 ■ Not available
 ■ No Information

Chapter 5

Maternal Mortality and Morbidity

5.1 Maternal mortality

The trends in maternal mortality over the last 10 years in the two districts compared to the national scenario are presented in figure 5.1.

The trends in all three districts show a decline. In Colombo and Gampaha the values appear to be

below the national level. High degree of fluctuation seen in Colombo and Kalutara districts are probably because of the small numbers each year.

Causes of death over this period of time has been examined for two 5 year time periods in tables 5.1(a), 5.1(b) and 5.1(c).

The trends in maternal mortality over the last 10 years in the two districts compared to the national scenario are presented in figure 5.1.




Table 5.1(a): Causes of maternal mortality by district 2001-2010 –Colombo

	2001	2002	2003	2004	2005	2001-2005	2006	2007	2008	2009	2010	2006-2010
MMR / 100,000 LB	14.9	40.4	19.2	38	39.1		33.9	35.9	30.97	43.3	13.0	
No. of all reported deaths	10	18	14	27	19	88	11	16	09	20	10	66
No. of confirmed maternal deaths	06	16	08	16	17	63	10	11	04	14	04	43
Type of maternal death												
Direct maternal deaths	05	07	07	10	08	37	04	07	00	11	04	26
Indirect maternal deaths	-	06	01	06	09	22	06	04	04	03	00	17
Inconclusive maternal deaths	01	03	-	-	-	04		0			00	00
Causes of maternal deaths												
Post partum haemorrhage	02		01	01	03	07	01	03		06		10
Pregnancy induced hypertension		03	03	02	02	10	01					01
Septic abortion	01	01	01	01		04		03				03
Cardiovascular disease		07	01	04	07	19	01	01	01	02		05
Pulmonary embolism	01	01	01			03	01					01
Amniotic fluid embolism						00				02	01	03
Suicide			01	02		03						00
Ectopic pregnancy				02	01	03						00
Liver disease	01			01		02		01		01		02
Renal Disease						00				01		01
Rupture of uterus		01				01						00
Cerebrovascular disease		01				01						00
Ante partum haemorrhage				01		01	01					01
Reproductive tract sepsis				01		01						00
Malignancy				01		01		01		01		02
Chicken pox					01	01						00
Medically complicated multiple pregnancy					01	01						00
Iatrogenic meningitis					01	01						00
Sepsis – Other					01	01	02		01		01	04
Pneumonia						00	01		02			03
Dengue Haemorrhagic fever						00	01	01				02
other						00	01	01				02
HELLP syndrome						00				01	01	02
Inconclusive	01	02				03						00
Anaesthetic drug related death						00					01	01

Source: National Maternal Mortality Review minutes

In the Colombo district over the two periods of time there has been a marked decrease in maternal deaths. There is decline in both direct and indirect obstetric deaths but the decline in deaths due to direct obstetric

causes is more marked. The number of deaths due to cardiovascular diseases have decreased to a third of the number in the earlier period but still constitutes the main contributor to indirect deaths.

Table 5.1(b): Causes of maternal mortality by district 2001-2010 -Gampaha

	2001	2002	2003	2004	2005	2001 - 2005	2006	2007	2008	2009	2010	2006-2010
MMR / 100,000 LB	36	44	22	24.4	60.6		35.3	34.7	25.5	21.7	20.3	
No. of all reported deaths	26	23	15	16	39	119	23	20	21	21	22	107
No. of confirmed maternal deaths	15	18	09	10	25	77	15	14	11	09	09	58
Type of maternal death												
Direct maternal deaths	13	13	09	09	15	59	08	07	07	05	02	29
Indirect maternal deaths	02	05	-	01	08	16	05	06	04	04	07	26
Inconclusive maternal deaths	-	-	-	-	02	02	02	01	00		00	03
Causes of maternal deaths												
Post partum haemorrhage	03	03	03	03	01	13	01	01	02	01		05
Pregnancy induced hypertension	02	02		01	04	09	03	02		01	02	08
Cardiovascular disease	01	02		01	05	09	02	02	01	03	03	11
Septic abortion	02	01				03	02	02		02	01	07
Pulmonary embolism	04	02	02	01		09		01				01
Reproductive tract sepsis		01	02	01	01	05			01			01
Liver disease		01	01		02	04	01	01	01			03
Ectopic pregnancy	02				02	04						00
Lower respiratory tract infection		01			01	02				01	01	02
Ante partum haemorrhage					01	01	01					01
Amniotic fluid embolism			01	01	01	03		01	01			02
Myocarditis		01			02	03	01				02	03
Anaesthetic death		01			01	02						00
Suicide (post partum psychosis)		01				01						00
Obstructed labour		01				01						00
Dengue fever		01				01						00
Deep vein thrombosis				01		01						00
Cerebrovascular disease				01		01						00
Aplastic anaemia					01	01						00
Renal disease					01	01						00
Iatrogenic meningitis					01	01						00
Acute disseminated encephalitis						00	01					01
Systemic lupus erythematosus						00	01		02			03
Sepsis – other					01	01						00
HELLP Syndrome						00			01	01		02
Other indirect	01					01						00
Inconclusive						00	02	01	01			04
Malignancy						00		03	01			04

Source: National Maternal Mortality Review minutes

In the Gampaha district the total number of maternal deaths has declined over the two five year periods of time. There has been a decrease in the deaths due to direct obstetric causes and an increase in the number

of deaths due to indirect causes. Cardiovascular disease is the main contributor to indirect deaths. Hypertensive disease, abortion and haemorrhage are the main contributors to direct obstetric deaths.

Table 5.1(c): Causes of maternal mortality by district 2001-2010 -Kalutara

	2001	2002	2003	2004	2005	2001-2005	2006	2007	2008	2009	2010	2006-2010
MMR / 100,000 LB	55.1	63	76	23	31		33.2	40.7	21.4	17.1	39.1	
No. of all reported deaths	14	21	21	18	18	92	20	15	11	11	18	75
No. of confirmed maternal deaths	12	13	12	05	07	49	07	07	05	04	09	32
Type of maternal death												
Direct maternal deaths	06	09	09	01	05	30	04	05	01	02	03	15
Indirect maternal deaths	06	04	03	04	01	18	03	01	04	02	06	16
Inconclusive maternal deaths	-	-	-	-	01	01	-	01	00	00	00	01
Causes of maternal deaths												
Cardiovascular disease	02	02		03	03	10	02	01	01		02	06
Pregnancy induced hypertension	01	04	05		02	12						00
Lower respiratory tract infection		02		01		03	01		01	01	01	04
Post partum haemorrhage		01	01		02	04			01			01
Pulmonary embolism	03	01				04						00
Ectopic pregnancy	01	01				02	01					01
Septic abortion	01		01			02						00
Myocarditis		01				01						00
Liver disease		01				01						00
Ante partum haemorrhage			01			01						00
Multiple sclerosis			01			01						00
Suicide			01			01					01	01
Cerebrovascular disease			01			01				01		01
Rupture of uterus						00	01				01	02
Reproductive tract sepsis						00	01					01
Multiple medical problems						00	01					01
Other indirect	04					04			01			01
Inconclusive			01	01		02		01				01
Amniotic fluid embolism						00		02	01	01		04
Diabetes Mellitus						00			01		01	02
HELLP syndrome						00				01		01
Sepsis other						00					02	02
Malignancy						00					01	01

Source: National Maternal Mortality Review minutes

5.2 Maternal morbidity

5.2.1 Admission patterns

Table 5.2: Admission patterns by type of institution

	Gampaha-Specialized		Gampaha-non Specialized		Kalutara – Specialized		Kalutara – non specialized		Colombo – Specialized		Colombo – Non Special		Western-		Western-Non Specialized	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Total BHT's Examined	2699		NI		1565		NI		3325		3		71		3	
Total BHT's reviewed in Obstetric wards	2408	89.2	NI	NI	1492	95.3	NI	NI	3033	91.2	1	33.3	6933	91.4	1	33.3
Outcome of admission																
Live discharge – (mother)	2392	99.3	NI	NI	1484	99.5	NI	NI	3007	99.1	1	100.0	6883	99.3	1	100.0
Transfer to other institution	9	0.4	NI	NI	6	0.4	NI	NI	22	0.7	0	0.0	37	0.5	0	0.0
Death	0	0.0	NI	NI	0	0.0	NI	NI	0	0.0	0	0.0	0	0.0	0	0.0
Needed critical care (ICU)	4	0.2	NI	NI	1	0.1	NI	NI	6	0.2	0	0.0	11	0.2	0	0.0
Transferred to Medical / Surgical unit	6	0.2	NI	NI	3	0.2	NI	NI	10	0.3	0	0.0	19	0.3	0	0.0
Reasons for admission																
Admitted for delivery	1661	69.0	NI	NI	884	59.2	NI	NI	1799	59.3	1	100.0	4344	62.7	1	100.0
Investigation of IUGR	52	2.2	NI	NI	47	3.2	NI	NI	66	2.2	0	0.0	165	2.4	0	0.0
Decrease foetal movements	98	4.1	NI	NI	18	1.2	NI	NI	142	4.7	0	0.0	258	3.7	0	0.0
Admitted for investigations	141	5.9	NI	NI	38	2.5	NI	NI	170	5.6	0	0.0	349	5.0	0	0.0
Febrile illness	8	0.3	NI	NI	7	0.5	NI	NI	22	0.7	0	0.0	37	0.5	0	0.0
Other illnesses	64	2.7	NI	NI	17	1.1	NI	NI	77	2.5	0	0.0	158	2.3	0	0.0
Other complications of pregnancy	170	7.1	NI	NI	147	9.9	NI	NI	396	13.1	0	0.0	713	10.3	0	0.0
Other	348	14.5	NI	NI	233	15.6	NI	NI	420	13.8	0	0.0	1001	14.4	0	0.0

It is seen that 3-6% of admissions in the Western province are for investigations. As could be expected in a district which has national referral centres for

EmONC nearly 13% of admissions in Colombo are for complications of pregnancy.

5.2.2 Delivery pattern

Table 5.3: Delivery patterns in specialist institutions and non-specialist institutions in Western province

	Gampaha-Specialized		Gampaha-non Specialized		Kalutara – Specialized		Kalutara – non specialized		Colombo – Specialized		Colombo – Non Special	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Total BHT's Examined	2699		NI		1565		NI		3325		3	
Total BHT's reviewed in Obstetric wards	2408	89.2	NI	NI	1492	95.3	NI	NI	2408	89.2	NI	NI
-												
Mode of delivery												
Vaginal delivery	1427	72.0	NI	NI	804	69.7	NI	NI	1526	65.9	0	0.0
Details documented	1427	100.0	NI	NI	804	100.0	NI	NI	1526	100.0	0	0.0
Cephalic	1374	96.3	NI	NI	785	97.6	NI	NI	1394	91.3	0	0.0
Breech	14	1.0	NI	NI	13	1.6	NI	NI	35	2.3	0	0.0
Vacuum	18	1.3	NI	NI	6	0.7	NI	NI	62	4.1	0	0.0
Forceps	21	1.5	NI	NI	0	0.0	NI	NI	35	2.3	0	0.0
CS - Elective	315	15.9	NI	NI	180	15.6	NI	NI	360	15.5	0	0.0
CS - Emergency	241	12.2	NI	NI	170	14.7	NI	NI	431	18.6	0	0.0
Total reported deliveries	1983	82.4	NI	NI	1154	77.3	NI	NI	2317	76.4	0	0.0

5.2.3 Complications of pregnancy and labour

Table 5.4 shows the morbidity pattern in specialised and non-specialised institutions. An attempt was made to collect data on abortion separately as

induced and spontaneous. However, only some institutions have reported the two separately, some reporting the two together as abortion. Therefore all data is presented as abortion.

Table 5.4 Complications of pregnancy and labour in specialist and non-specialist institutions in Western province

	Gampaha-Specialized		Gampaha-non Specialized		Kalutara – Specialized		Kalutara – non specialized		Colombo – Specialized		Colombo – Non Special		Western-Specialized		Western – Non Specialized	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Total BHT's Examined	2699		NI		1565		NI		3325		3		7589		3	
Complications of Labor																
Prolonged / Obstructed labor	35	1.3	NI	NI	12	0.8	NI	NI	47	1.4	0	0.0	94	1.2	0	0
Lack of progress	61	2.3	NI	NI	29	1.9	NI	NI	87	2.6	0	0.0	177	2.3	0	0
Retained placenta	13	0.5	NI	NI	8	0.5	NI	NI	28	0.8	0	0.0	49	0.6	0	0
Tears	49	1.8	NI	NI	7	0.4	NI	NI	78	2.3	0	0.0	134	1.8	0	0
Cord prolapse	4	0.1	NI	NI	0	0.0	NI	NI	6	0.2	0	0.0	10	0.1	0	0
Shoulder dystocia	2	0.1	NI	NI	0	0.0	NI	NI	4	0.1	0	0.0	6	0.1	0	0
Other	39	1.4	NI	NI	4	0.3	NI	NI	81	2.4	0	0.0	124	1.6	0	0
Total	203	7.5	NI	NI	60	3.8	NI	NI	331	10.0	0	0.0	594	7.8	0	0
Hemorrhage																
Placenta previa	13	0.5	NI	NI	7	0.4	NI	NI	13	0.4	0	0.0	33	0.4	0	0
Accreta/increta/percreta placenta	2	0.1	NI	NI	0	0.0	NI	NI	0	0.0	0	0.0	2	0.0	0	0
Abruptio placenta	9	0.3	NI	NI	1	0.1	NI	NI	6	0.2	0	0.0	16	0.2	0	0
Ruptured uterus	1	0.0	NI	NI	0	0.0	NI	NI	0	0.0	0	0.0	1	0.0	0	0
Postpartum hemorrhage	16	0.6	NI	NI	8	0.5	NI	NI	17	0.5	0	0.0	41	0.5	0	0
Other obstetric hemorrhage	14	0.5	NI	NI	1	0.1	NI	NI	2	0.1	0	0.0	17	0.2	0	0
Total	55	2.0	NI	NI	17	1.1	NI	NI	38	1.1	0	0.0	110	1.4	0	0
Infection																
Post-Partum sepsis	11	0.4	NI	NI	4	0.3	NI	NI	6	0.2	0	0.0	21	0.3	0	0
Pyelonephritis	5	0.2	NI	NI	0	0.0	NI	NI	12	0.4	0	0.0	17	0.2	0	0
Influenza-like illness (viral fevers)	9	0.3	NI	NI	2	0.1	NI	NI	5	0.2	0	0.0	16	0.2	0	0
Other systemic infections/sepsis	24	0.9	NI	NI	7	0.4	NI	NI	20	0.6	0	0.0	51	0.7	0	0
Total	49	1.8	NI	NI	14	0.9	NI	NI	44	1.3	0	0.0	107	1.4	0	0
Hypertension																
Chronic hypertension	10	0.4	NI	NI	4	0.3	NI	NI	19	0.6	0	0.0	33	0.4	0	0
PIH other than eclampsia	102	3.8	NI	NI	51	3.3	NI	NI	218	6.6	0	0.0	371	4.9	0	0
Eclampsia	7	0.3	NI	NI	4	0.3	NI	NI	18	0.5	0	0.0	29	0.4	0	0
Total	119	4.4	NI	NI	59	3.8	NI	NI	255	7.7	0	0.0	433	5.7	0	0
Total gynaecological tickets examined	291	10.8	NI	NI	73	4.7	NI	NI	292	8.8	2	66.7	656	8.6	2	66
Gynaecological conditions																
Ectopic pregnancies	22	0.8	NI	NI	2	0.1	NI	NI	19	0.6	0	0.0	43	0.6	0	0
Abortions	260	9.6	NI	NI	71	4.5	NI	NI	272	8.2	2	66.7	603	7.9	2	66
Abortion related hemorrhage	14	0.5	NI	NI	3	0.2	NI	NI	3	0.1	0	0.0	20	0.3	0	0
Abortion related sepsis	9	0.3	NI	NI	0	0.0	NI	NI	2	0.1	0	0.0	11	0.1	0	0
Known Induced	9	0.3	NI	NI	3	0.2	NI	NI	1	0.0	0	0.0	13	0.2	0	0
Other abortions	260	9.6	NI	NI	68	4.3	NI	NI	272	8.2	2	66.7	600	7.9	2	66

Nearly 6% of admissions to specialised units in the province are for hypertensive disease in pregnancy. The other obstetric complications are comparatively low. Known induced abortions, abortion related haemorrhage, sepsis are also low.

Data from non specialised units are available only in the Colombo district. Complications are not seen in the hospitals without specialised services.

5.2.4 Co-morbidities

As could be expected the co-morbidity rates are comparatively high in the Colombo district (10%) the main condition being gestational diabetes, diabetes prior to pregnancy and impaired glucose tolerance taken together.

Table 5.5 Co-morbidities in specialist institutions and non-specialist institutions in Western province.

	Gampaha-Specialized		Gampaha-non Specialized		Kalutara – Specialized		Kalutara – non specialized		Colombo – Specialized		Colombo – Non Special		Western-Specialized	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No	%
Total BHT's Examined	2699		NI		1565		NI		3325		3		7589	
Co Morbidities														
Severe Anemia	13	0.5	NI	NI	0	0.0	NI	NI	13	0.4	0	0.0	26	0.3
Heart disease	30	1.1	NI	NI	15	1.0	NI	NI	55	1.7	0	0.0	100	1.3
Lung disease	33	1.2	NI	NI	14	0.9	NI	NI	36	1.1	0	0.0	83	1.1
Renal disease	5	0.2	NI	NI	2	0.1	NI	NI	5	0.2	0	0.0	12	0.2
Hepatic disease	1	0.0	NI	NI	0	0.0	NI	NI	6	0.2	0	0.0	7	0.1
Gestational diabetes	53	2.0	NI	NI	24	1.5	NI	NI	170	5.1	0	0.0	247	3.3
Diabetes prior to pregnancy	8	0.3	NI	NI	13	0.8	NI	NI	19	0.6	0	0.0	40	0.5
Impaired Glucose Tolerance	14	0.5	NI	NI	9	0.6	NI	NI	12	0.4	0	0.0	35	0.5
Total	157	5.8	NI	NI	77	4.9	NI	NI	316	9.5	0	0.0	550	7.2

5.3 Transfers


Table 5.6: Pattern of transfers in the Western province

Hospital	Usual place of emergency transfer	Average time taken for a transfer in minutes	Number of obstetric cases transferred during the prospective survey (4 weeks)	Number of transfers with documented direct obstetric complications
Gampaha District				
TH Ragama	NI	NI	1	0
DGH Gampaha	DMH	45 minutes	0	0
DGH Negombo	TH Ragama	30 minutes	8	0
BH Wathupitiwala	DMH	90 minutes	0	0
BH Mirigama	BH Wathupitiwala	40 minutes	NI	NI
BH Kiribathgoda	TH Ragama	30 minutes	NI	NI
DH Minuwangoda	DGH Gampaha	15 minutes	NI	NI
Colombo District				
TH CSHW	NI	NI	0	0
TH DMH	NI	NI	3	0
CSTH	NI	NI	2	1
SJGH	NI	NI	NI	NI
BH Awissawella	DMH	45 minutes	5	2
BH Homagama	CSHW	45 minutes	2	0
BH Mulleriyawa	CSTH	20 minutes	14	4
DH Piliyandhala	CSTH	30 minutes	0	0
Kalutara District				
GH Kalutara	NI	NI	3	0
BH Horana	CSTH	45 minutes	2	0
KMH	CSTH	45 minutes	1	1
BH Pimbura	GH Kalutara	25 minutes	NI	NI
DH Mathugama	GH Kalutara	20 minutes	NI	NI
DH Beruwala	GH Kalutara	30 minutes	NI	NI

NI (No information available from the hospital)

The transfer times within the district and the province are low.

Figure 5.2 Pattern of transfers in the Western province



Chapter 6

Resources for EmONC - Infrastructure

Institutional services for pregnant women are provided through a graded network of 603 hospitals spread throughout the country which have specially identified maternity wards. The infrastructure of these institutions has a significant influence on the provision of quality obstetric and neonatal care. This chapter summarizes the infrastructure resource distribution in the country at the time of survey. The source of information used for each table is indicated. Many of the tables are self-explanatory.

In the absence of norms/standards on infrastructure based on services, workload and type of institution the findings of this survey are presented so that they may be used as a guide for developing these nationally.

6.1 Facility profile

This section describes the facility profile by district and institution. The information presented in table 6.1 was obtained from the heads of the institutions.

Table 6.1 Summary of services provided by institution

	Colombo								Gampaha								KALUTARA						
	TH CSHW	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH COLOMBO NORTH	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		DGH KALUTARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA	
Facilities available for maternal care																							
Bed strength of the hospital	494	343	1093	495	426	285	95		1405	699	657	564	148	84	116		878	370	442	137	112	65	
No. of obstetrics units available	5	5	3	2	2	1	1		3	2	2	2	1	0	1		2	2	2	1	1	1	
No. of labor rooms available	5	5	2	1	1	1	1		2	2	1	1	1	0	1		1	1	1	1	1	1	
Total beds dedicated for obstetric care	304	230	NI	100	63		13		NI	NI	97	131	10	6	11		56	NI	62	11	12	7	
Proportion of Obs beds / total beds of the institution	61.5	67.1		20.2	14.8	0.0	13.7				14.8	23.2	6.8	7.1	9.5		6.4		14.0	8.0	10.7	10.8	
Beds dedicated for obs care - antenatal	135	98		50	30	13	9				56	80	5	5	5				38	6	8	5	
Beds dedicated for obs care - postnatal	169	132		50	33	9	4				41	51	5	0	6				24	5	4	2	
NICU	✓	✓	✓	✗	✗	✗	✗		✗	✓	✗	✗	✗	✗	✗		✗	✗	✗	✗	✗	✗	
SCBU	✓	✓	✓	✗	✗	✗	✗		✓	✓	✓	✓	✗	✗	✗		✓	✓	✓	✗	✗	✗	
MBC	✓	✗	✗	✗	✗	✗	✗		✗	✓	✗	✗	✗	✗	✗		✗	✓	✗	✗	✗	✗	
LMC	✓	✓	✓	✓	✗	✗	✗		✓	✓	✗	✓	✗	✗	✗		✓	✓	✓	✓	✗	✗	
Antenatal clinic care	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	
Post natal clinic care	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	
Cervical screening	✓	✓	✓	✗	✓	✓	✗		✓	✗	✓	✓	✗	✓	✗		✗	✗	✗	✓	✗	✓	
Family planning clinic	✓	✓	✓	✓	✓	✓	✓		✓	✓	✗	✓	✓	✗	✓		✗	✓	✓	✓	✓	✓	
Well baby clinic	✓	✓	✓	✓	✓	✓	✓		✓	✓	✗	✓	✓	✓	✗		✓	✓	✓	✓	✓	✓	
Health education unit	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✗	
Infection control unit	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✗	✗	

Table 6.2 provides a summary of the institutional profiles in terms of facilities necessary for the provision of EmONC services. The criteria used in the classification of EmONC services have been described in chapter 2. Information necessary were obtained from the heads of institutions and personnel in charge of obstetric wards, labour rooms and operating theatres.

6.2. Labour room

Tables 6.3 (a) to 6.3 (c) provides a summary of the labour room infrastructure facilities available by type of hospital. Data on labour room infrastructure facilities was collected based on the requirements listed in the Labour Room Management Guide (FHB, 2007). It is expected that each of the items listed in column one be available in all institutions.

Table 6.2: Summary of institutional profiles in districts and the province

Hospital	Western Province																	
	Gampaha								Colombo								Kaltara	
	TH RAGAMA	DGH GAMPABA	DGH NEGOMBO	DH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		TH CSHW	TH DMH	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		GH Kaltara	Kethumathie MH
Comprehensive EmONC status	+	+	+	+	-	-	-		+	+	+	+	+	-	-		+	+
Basic EmONC status	-	-	-	-	-	-	-		-	-	-	-	-	-	-		-	-
24*7 Comprehensive EmONC status	+	+	+	+					+	+	+	+	+				+	+
Institutions with 24*7 electricity	+	+	+	+	-	-	+		+	+	+	+	+	+	-		+	+
Institutions with 24*7 water supply	-	+	+	+	+	-	+		+	+	-	+	+	+	+		+	-
Institutions with 24*7 Transport	+	+	-	-	+	-	-		+	+	-	+	+	-	-		+	-
Institutions with 24*7 Communication	+	+	+	+	+	+	-		+	+	+	+	+	+	+		+	+
Availability of a functional theatre	+	+	+	+	-	-			+	+	+	+	+	+			+	+
Availability of a functional blood bank	+	+	+	+	-	-			+	+	+	+	+	+			+	+
24*7 obstetric surgical cover	+	+	+	+	-	-			+	+	+	+	+	+			+	+
24*7 Anesthetic cover	+	+	+	+	-	-			+	+	+	+	+	+			+	+
24*7 Blood transfusion service cover	+	+	+	+	-	-			+	+	+	+	+	+			+	+
Dedicated Obstetric theatre	-	-	-	-					+	+	+	+	+	-	-		+	+

Table 6.3 (a) Summary of labour room facilities by institution – Infrastructure

INSTITUTION	COLOMBO																GAMPAHA										KALUTARA									
	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH DMH	TH DMH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERYAWA	DH PILIYANDALA	TH RAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH GAMPAHA	DGH NEGOMBO	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTARA	DGH KALUTARA	BH PANADURA	BH HOREANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA	
UNIT	1	2	3	4	5	7	15	3	5	16	14	18	21	1+2	1	2			17	18	25	10	12	10	12			NF		A	B					
Water Supply																																				
Water supply	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Elbow or foot operated tap for hand washing	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Electricity and Illumination																																				
Generator / separate generator supply	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Emergency lamps	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
At each head-end of bed three, 13A plug points	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
At resuscitation area three, 13A plug points	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
One, 15A plug point	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Communication																																				
Cordless telephone for PHO	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Direct telephone line to labour room	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Intercom facilities	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Alarm for emergencies	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Ventilation																																				
AC the temperature maintained at 25°C	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Exhaust fan	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Presence of windows above 1.2m	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

Table 6.3(b) Summary of labour room facilities by institution – infrastructure

INSTITUTION	COLOMBO																GAMPAHA										KALUTARA											
	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH DMH	TH DMH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH HOMAGAMA	BH MULLERIVAWA	DH PILIYANDALA	TH RAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH GAMPAHA	DGH NEGOMBO	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTARA	DGH KALUTARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA		
UNIT	1	2	3	4	5	7	15	3	5	16	14	18	21	1+2	1	2			17	18	25	10	12	10	12				NP		A	B						
Walls and Floor																																						
Walls tiled up to 1.8m from floor																																						
Tiled floor																																						
At the entrance to labour room																																						
Place to change clothes																																						
Place for hand washing																																						
Shoe rack																																						
Automatically closing doors to labour room																																						
Washroom																																						
Separate room for washing instruments																																						
Steel/fiberglass sink																																						
Racks																																						
Two 13A plug points																																						
Separate entrance from outside																																						
Toilet Facilities																																						
Separate toilets for labour room																																						
Commode type of toilets																																						
Squatting type of toilets																																						
Railings on either side of wall (for squatting)																																						
Other requirements																																						
Separate trolley for labour room																																						
Place for temporary storage of dirty linen etc																																						
Use of colour card system for waste disposal																																						
Notice board outside the labour room																																						
On call doctor's information displayed in LR																																						

Table 6.3(c) Summary of labour room facilities by institution - infrastructure

INSTITUTION	COLOMBO																	GAMPAHA										KALUTARA								
	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH DMH	TH DMH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH HOMAGAMA	BH MULLERIYAWA	DH PILYANDALA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH GAMPAHA	DGH NEGOMBO	DGH NEGOMBO	BH WATHUPITTIALA	BH MTRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTARA	DGH KALUTARA	BH PANADURA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA		
UNIT	1	2	3	4	5	7	15	3	5	16	14	18	21	1+2	1	2			17	18	25	10	12	10	12			NE		A	B					
Human Resources																																				
Nurses-midwifery trained	✓	✓	✗	✓	✓			✓	✓	✓					✗		✓	✓	✓						✓				✓	✓		✓	✓	✓	✓	
Nurses-without midwifery training		✗			✗				✗		✗		✗	✗			✗								✗	✗			✓	✓	✗	✗	✓	✓	✓	
Midwives	✓	✓	✓	✓				✗	✗	✓							✗			✓						✓	✓			✓	✓	✓	✓	✓	✓	
24hr on-call house officer	✓	✓		✓			✗			✓						✓								✓	✓	✓	✗			✗		✓	✓	✓	✗	
Newborn care area																																				
Resuscitator	✓	✓	✓					✗		✓								✗									✗			✗						
Neonatal cots	✓	✓					✗		✓	✗							✗									✗	✗			✓	✓		✗	✓	✓	
Sink with water supply	✓	✗	✗	✗	✗	✗				✗		✗	✗	✗			✗			✓						✗	✗			✓	✓	✓	✓	✓	✓	
Low tech incubators	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗			✗	✗		✓				✗	✗	✗	✓			✗	✗	✗	✗	✗	✗	
Furniture and General Items																																				
Delivery beds with stirrups	✓	✓	✓	✓	✓					✓						✗			✓					✗	✗	✓	✓			✓	✓	✓	✓	✓	✓	
Spot lamps (movable)	✓	✓	✓	✓	✓			✓	✓	✓								✓						✓	✓	✓	✓			✓	✓	✓	✓	✓	✓	
Wall clock																																				
Oxygen cylinder stands and regulators																																				
Refrigerators	✓	✓	✓		✗		✗	✗																			✗			✓	✓		✗	✓	✗	
Neonatal suckers	✓	✓	✓	✓		✗	✗	✗	✗							✗		✓						✗	✗	✓			✓	✓	✓	✓	✓	✗	✗	
Equipments																																				
Delivery sets	✓	✓								✓																✓	✓									
Pinard foetal stethoscope	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓					✗		✓		✓						✓	✓	✓			✓	✓	✓	✓	✓	✓	
Doppler (Foetal heart detector)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓								✗		✓						✓			✓	✓	✓	✓	✓	✓	✓	
CTG machine	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓								✗		✓				✗	✗	✓	✓			✗				✗	✗	
Vacuum extractor													✗					✗					✗	✗	✗	✓				✓	✓	✓	✓	✓	✓	
Wrigley's forceps							✗	✓										✗		✗									✓	✓			✓	✓	✓	
Secca weighing scale								✗				✗		✗				✗	✗	✗				✗		✓							✓	✓	✓	
Infant Magills laryngoscope(straight blade)	✓	✓	✓	✓						✗								✗	✗	✗	✗			✗		✓				✓	✓	✓	✓	✓	✓	
Wall thermometer	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗		✗	✗	✗	✗	✗	✗	✗	
Surgical consumables																																				
Cord clamps/Sterile threads	✓	✓	✓	✓		✗	✗	✗	✓							✓													✓	✓		✓	✓	✓	✓	
Neonatal ambu bag	✓	✓							✓	✓						✓			✓							✓			✓	✓	✓	✓	✓	✓	✓	
Neonatal sucker tubes	✓	✓					✗											✗		✓						✓			✓	✓	✓	✓	✓	✓	✗	
Naso-gastric tubes	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓									✓						✓				✓	✓			✓		✓	
Endo tracheal tubes																			✓						✓	✓								✓	✓	
Size 2.5	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓								✗	✗							✗	✗			✗			✗	✗	✗	
Size 3	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓								✗	✗										✗			✗	✗	✗	✗	
Size 3.5	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓								✗	✗										✗			✗	✗	✗	✗	
Size 4	✗	✗	✗	✗	✗	✗	✗	✓				✗	✗	✗	✗			✗	✗													✗	✗	✗	✗	
Flexible stiletto(for ET tube)	✓	✓	✓	✓						✗		✗	✗	✗	✗			✗	✗					✗		✗	✗		✗	✓	✓	✓	✓	✓	✓	
IV cannula	✓	✓	✓	✓						✓								✓											✓	✓					✓	✓
Disposable syringes																				✗									✓						✓	
Drugs																																				
Adrenaline	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓									✓											✓	✓	✓	✓	✓	✓	
Naloxone																			✗											✓	✓	✓	✓	✗	✗	✗
Vitamin K																			✗											✓			✗	✗	✗	
Normal Saline																														✓						

6.3. Operating Theatre

A functioning operating theatre is an absolutely essential facility to provide comprehensive emergency obstetric care. The information on the theatre facilities were obtained from the officer in charge of the theatre in each institution and are presented in table 6.4.

6.4. Blood transfusion services

The ability to provide blood transfusion is one of the signal functions that differentiate between BEmONC and CEmONC services. It is a lifesaving intervention in an emergency and timely and safe blood transfusion services should be available 24*7 every day of the year. The information on blood transfusion services were obtained from the medical officer- in charge of the blood bank in each institution.

Table 6.4: Operation theatre facilities by institution

	COLOMBO						GAMPAHA						KALUTHARA			
	TH CSHW	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	TH COLOMBO NORTH	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA
Theatre complex	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✗	✓	✓	✓	✗
Functioning or not	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Separate operating theatre only for obstetric patients	✓	✓	✓	✓	✗	✗	✗	✗	✗	✗	✗	✗	✗	✓	✗	✗
Arrangement during working hours																
Interrupt the Surgical list					✓				✓	✓			✓		✓	
Interrupt the Gynae list					✓	✓	✓	✓	✓	✓			✓		✓	
Arrangement during off hours																
Theatre is freely available					✓	✓										
disturb a casualty list .							✓	✓	✓	✓			✓		✓	
During the past two months a delay of > 20 mins from the time of request due to the OT being occupied by another surgery	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗			✗	✗	✗	
Equipments and Supplies																
Availability of an operating table	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓			✓	✓	✓	
In functional condition	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓			✓	✓	✓	
Availability of adjustable light	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓			✓	✓	✓	
The light is shadow less	✓	✓	✓	✗	✓	✗	✓	✓	✓	✓			✓	✓	✓	
Requested for additional linen during the past three months	✗	✓	✓	✓	✓	✗	✓	✓	✓	✓			✗	✗	✗	
Use separate packs (laparotomy packs, LSCS packs) for surgeries	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓			✓	✓	✓	
Number of LSCS packs	10	12	7	6	5	3	11	8	8	7			8	5	5	
Number of laparotomy packs	6	4	2	4	2	1	6	3	2	2			2	1	1	
Postponing emergency obs surgery	✗	✓	✗	✗	✗	✗	✗	✗	✗	✗			✗	✗	✗	
Lack of staff	✗	✗	✗	✗	✗		✗	✗	✗	✗						
Lack of equipment	✗	✗	✗	✗	✗		✗	✗	✗	✗						
Lack of water and electricity facilities	✗	✗	✗	✗	✗		✗	✗	✗	✗						
Lack of linen	✗	✗	✗	✗	✗		✗	✗	✗	✗						
Lack on anaesthetic drugs or gases	✗	✗	✗	✗	✗		✗	✗	✗	✗						
Malfunctioning of equipments, A/C	✗	✓	✗	✗	✗		✗	✗	✗	✗						
Infrastructure																
Days without electricity	0	1	0	0	0	0	1	0	0	0			0	0	0	
Separate generator for the theatre	✗	✗	✓	✗	✓	✗	✗	✗	✗	✓			✗	✗	✗	
Dedicated staff for generator	✗	✓	✓	✓	✓	✓		✗	✗	✗			✓	✓	✓	
Continuous supply of water	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓			✓	✓	✓	
Days without water	✗	✗	✗	✗	✗	✗	✓	✗	✗	✗			✗	✗	✗	
Direct telephone line to the theatre	✓	✗	✓	✗	✓	✓	✗	✓	✓	✓			✓	✓	✓	
Direct dialing facility in the theatre	✓	✗	✗	✗	✗	✗	✗	✗	✗	✗			✗	✗	✗	
Cell phones	✓	✗	✓	✓	✓	✓	✓	✓	✗	✓			✓	✓	✓	
Resuscitator at the newborn corner of the OT	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓			✓	✓	✓	

Table 6.5: Blood transfusion services by institution

	COLOMBO							GAMPAHA							KALUTARA			
	TH CSHW	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA		TH COLOMBO NORTH	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITIWALA	BH MIRIGAMA	BH KIRIBATHGODA		DGH KALUTARA	BH PANADURA	BH HORANA	BH PIMBURA
Availability of a blood bank	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✗	✗		✓	✓	✓	✗
Mismatch transfusions recorded during the past six months	✓	✗	✗	✗	✗	✗		✗	!	✗	✗				✗	✗	✗	
Instances of septicemia following blood transfusions recorded during the past six months	✗	✗	✗	✗	✗	✗		✗	✗	✗	✗				✗	✗	✗	
Availability of blood products																		
Whole blood	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓				✓	✓	✓	
Platelets	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓				✓	✓	✓	
FFP	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓				✓	✓	✓	
Cryoprecipitate	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓				✓	✓	✓	
O negative blood in stock	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓				✓	✓	✓	
O positive blood in stock	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓				✓	✓	✓	
Cross matching of blood during normal working hours																		
MO/Blood Bank	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓				✓	✓	✓	
House Officer																		
Cross matching of blood after normal working hours																		
MO/Blood Bank	✓	✓	✓	✗	✗	✓		✓	✗	✓	✗				✓		✓	
House Officer				✓	✓			✓	✓	✓	✓				✓	✓		

6.5. Laboratory services

Following information on laboratory was obtained from the Medical officer- in charge of the laboratory or chief MLT.

Table 6.6: Laboratory services by institution

	COLOMBO								GAMPAHA								KALUTHARA					
	TH CSHW	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH COLOMBO NORTH	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA
A functioning laboratory is available	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✗	✓		✓	✓	✓	✗	✓	✓
Laboratory facilities available for 24*7	✗	✓				✗	✗							✗	✗		✗	✗	✗	✗	✗	✗
The operational arrangement of maintaining the 24*7 service		OC	OC	OC	OC	OC			R	R	OC	R					OC		OC			
Testing facilities available																						
DURING NORMAL WORKING HOURS																						
a. Haemoglobin	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✗
b. WBC/DC	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
c. Platelets	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
d. PCV	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✗	✗
e. Blood urea	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✗	✗
f. Blood sugar	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✗	✗
g. Bilirubin	✓	✓	✓	✗	✓	✓	✗		✓	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✗	✗
h. Grouping & Rh	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✗	✗
i. Electrolytes	✓	✓	✗	✓	✓	✓	✗		✓	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✗	✗
j. Blood culture & ABST	✓	✓	✗	✓	✓	✓	✗		✓	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✗	✗
k. Urine culture & ABST	✓	✓	✗	✓	✓	✓	✗		✓	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✗	✗
l. BT / CT	✓	✓	✗	✓	✓	✓	✗		✗	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✗	✗
m. Urine bile	✓	✓	✗	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
n. ECG	✓	✓	✗	✓	✓	✓	✓		✗	✓	✗	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
o. X ray	✓	✓	✓	✓	✓	✓			✗	✓	✗	✓	✗		✗		✓	✓	✓	✗	✗	✗
AFTER NORMAL WORKING HOURS																						
a. Haemoglobin	✓	✓	✓	✓	NI				✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
b. WBC/DC	✓	✓	✓	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
c. Platelets	✓	✓	✓	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
d. PCV	✓	✓	✓	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
e. Blood urea	✓	✓	✓	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
f. Blood sugar	✓	✓	✓	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
g. Bilirubin	✓	✓	✓	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
h. Grouping & Rh	✓	✓	✓	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
i. Electrolytes	✓	✓	✗	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
j. Blood culture & ABST	✗	✓	✗	✗					✗	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
k. Urine culture & ABST	✗	✓	✗	✗					✗	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
l. BT / CT	✓	✓	✗	✓					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
m. Urine bile	✓	✓	✗	✗					✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
n. ECG	✓	✓	✓	✓					✓	✓	✗	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
o. X ray	✓	✓	✓	✓					✓	✓	✗	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓

6.6: Infection Control

The information in table 6.7(a) was collected from the in charge of the CSSD and table 6.7(b) was collected from the sister or nursing officer in charge of labour rooms and the inquiry refers to the day of collection of data.

6.7: Pharmacy and Medicines for EmONC services

The information in tables 6.8a, 6.8b and 6.9 was collected from the officer in-charge of the drug stores of the institution and the inquiry refers to the day of collection of data.

Table 6.7 (a): Availability of infection control measures by institution

	COLOMBO								GAMPAHA								KALUTHARA					
	TH CSHW	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH COLOMBO NORTH	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA
Availability of a separate CSSD	✓	✓	✓	✓	✓	✗	✗		✓	✓	✗	✓	✗	✗	✗		NI	✗	✗	✗	✗	✗
If no CSSD:																						
Separate autoclave room						✗	✗								✓					✗	✗	
Autoclave (with temperature and pressure gauges)							✓					✓		✓			✓	✗			✓	
Hot air Sterilizer (dry oven)						✗	✗											✗		✗	✗	
Steam Sterilizer						✗	✗				✓				✓			✗		✗	✓	
Sterilizer / Pressure Cooker (electric)						✗	✓				✗				✗			✗		✓	✗	
Sterilizer / Pressure Cooker (kerosene heated)						✓	✓				✗				✗			✗		✗	✗	
Sterilization drum						✗	✓				✓				✓			✓		✓	✓	
Sterilization drum stand						✗					✗				✓			✗		✗	✓	
Availability of a policy on waste disposal	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✗	✓		✓							
It is functioning	✓	✓	✓	✓	✓	✓			✓	✓	✓		✗		✓							
Availability of a specific methodology of biological waster such as placenta	✓	✓	✓	✓		✓			✓	✓	✓	✓	✗		✗		✓	✗		✗	✓	
If yes describe briefly																						
PIT												✓										
Buried			✓	✓		✓				✓					✓						✓	
incinerate											✓						✓					
Refridgerate and cremation	✓	✓							✓													
Availability of a functioning incinerator	✗	✗	✗	✗	✗	✓			✓	✓	✗	✗	✗		✗							

Table 6.7 (b): Infection control measures by institution

	COLOMBO																GAMPAHA										KALUTHARA								
	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH DMH	TH DMH	TH DMH	TH DMH	TH COLOMBO SOUTH	TH COLOMBO SOUTH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA	TH COLOMBO NORTH	TH COLOMBO NORTH	TH COLOMBO NORTH	DGH GAMPAHA	DGH GAMPAHA	DGH NEGOMBO	DGH NEGOMBO	BH WATHUPITIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA	
INFECTION PREVENTION																																			
Soap	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	NI	✓	✓	✓	✓	✓	✓	✗
Antiseptics	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓
Gloves	✓	✓	✓	✓	✓	✗	✗	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Heavy duty gloves	✓	✓	✗	✗	✓	✗	✓	✓	✗	✓	✓	✗	✓	✗	✗	✓	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Non-sterile protective clothing	✓	✓	✓	✓	✓	✗	✗	✗	✗	✓	✓	✗	✓	✗	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Bleach or bleaching powder	✓	✓	✓	✓	✓	✗	✓	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Prepared disinfection solution	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Covered contaminated waste trash bin	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Puncture proof sharps container	✓	✓	✓	✓	✓	✗	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Protective equipment for HIV	✓	✓	✓	✓	✓	✗	✗	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Chlorhexidine	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Ethanol (70%)	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Polyvidone iodine	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Hand rub	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

Table 6.8 (a): Availability of essential drugs by institution

	COLOMBO								GAMPAHA								KALUTHARA					
	TH CSHW	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH COLOMBO NORTH	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA
ANTIBIOTICS																						
Amoxicillin	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Ampicillin	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✗	✓	✓
Benzyl Penicillin	✓	✓	✓	✓	✓	✓	✓		✓	✗	✓	✓	✓	✓	✓		✓	✓	✓	✗	✓	✓
Cefalexin	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✗	✓	✓
Cefotaxime	✓	✓	✓	✓	✓	✓	✗		✓	✗	✓	✓	✗	✓	✗		✓	✓	✓	✗	✗	✗
Ceftriaxone	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✗	✓	✗		✓	✓	✓	✗	✗	✗
Cefuroxime	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✗	✓	✗
Cloxacillin	✓	✓	✓	✓	✓	✓	✓		✗	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Phenoxymethyl penicillin	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✗	✓	✓	✓		✓	✓	✓	✗	✓	✓
Erythromycin	✓	✓	✓	✓	✓	✓	✓		✗	✗	✓	✓	✗	✗	✓		✓	✓	✓	✗	✓	✓
Gentamicin	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Metronidazole	✓	✓	✓	✓	✓	✓	✓		✓	✗	✓	✓	✓	✓	✓		✓	✗	✓	✓	✓	✓
ANTICONVULSANTS																						
Magnesium sulphate (injection)	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✗	✓	✓		✓	✓	✓	✗	✓	✓
Diazepam (injection)	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✗	✓	✓	✓	✓
Phenobarbital	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Phenytoin	✓	✓	✓	✓	✓	✓	✓		✗	✓	✓	✗	✗	✗	✓		✓	✓	✓	✓	✓	✓
ANTIHYPERTENSIVES																						
Hydralazine	✓	✓	✗	✓	✓	✓	✓		✓	✓	✓	✓	✗	✗	✗		✓	✓	✓	✗	✗	✓
Labetalol	✓	✗	✗	✗			✗		✗	✗	✗	✗	✗	✗	✗		✗	✗	✗	✗	✗	✗
Methyldopa	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✗	✓	✓	✓		✓	✓	✓	✓	✓	✗
Nifedipine	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
OXYTOCICS																						
Ergometrine	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✗	✓		✓	✓	✓	✓	✓	✓
Oxytocin	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✗	✓	✓		✓	✓	✓	✓	✓	✓
Oxytocin + Ergometrine	✗	✗	✗	✗			✗		✗	✗	✗	✗	✗	✗	✗		✗	✗	✗	✗	✗	✗
EMERGENCY DRUGS																						
Adrenaline (Epinephrine)	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Atropine sulphate	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Calcium gluconate	✓	✓	✓	✓	✓	✓	✗		✗	✗	✓	✓	✓	✗	✗		✓	✓	✓	✗	✗	✗
Digoxin	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✗	✓	✓	✓		✓	✗	✓	✓	✓	✓
Ephedrine	✓	✓	✓	✓	✓	✓	✓		✗	✗	✓	✓	✗	✗	✗		✓	✓	✓	✗	✗	✓
Frusemide	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Hydrocortisone	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Naloxone	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✗	✗	✗		✓	✓	✓	✗	✗	✗
Promethazine	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
ANALGESICS																						
Morphine	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Pethidine	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
TOCOLITICS																						
Nifidipine	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Salbutamol	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✗	✓	✓	✓		✓	✓	✓	✓	✓	✓

Table 6.8 (b): Availability of essential drugs by institution

	COLOMBO								GAMPAHA								KALUTHARA					
	TH CSHW	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH COLOMBO NORTH	DGH GAMPAA	DGH NEGOMBO	BH WATHUPITIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA
STEROIDS																						
Betamethasone	✓	✓	✓				✗		✓	✗	✗	✗	✗	✗	✓		✓	✗	✗	✗	✓	✗
Dexamethasone	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Prednisolone	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
IV FLUIDS																						
Dextrose	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Normal saline	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Ringer's lactate	✓	✓	✗	✓	✓	✓	✗		✓	✓	✓	✗	✓	✓	✗		✓	✓	✓	✓	✗	✗
Starch	✓	✓	✓	✓			✗		✓	✓	✓	✓	✗	✓	✗		✓	✗	✗	✗	✗	✗
ANTIMALARIALS																						
Chloroquine	✗	✗	✓	✓			✗		✓	✓	✓	✓	✗	✗	✗		✓	✗	✗	✓	✗	✗
Quinine Dihydrochloride	✗	✗	✓	✗			✗		✓	✗	✗	✓	✗	✗	✗		✗	✗	✗	✗	✗	✗
ANTIRETROVIRALS	Available at the district STD clinic																					
CONTRACEPTIVES																						
Pills	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✗		✗	✓	✓	✓	✗	✓
Condoms	✓	✓	✓	✓	✓	✓	✓		✓	✗	✓	✓	✓	✗	✓		✗	✓	✓	✓	✗	✓
DMPA	✓	✓		✗			✓		✓	✗	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
IUD	✓	✓		✓			✓		✓	✗	✓	✓	✓	✓	✓		✗	✓	✓	✓	✓	✓
Implants	✓	✓		✗			✗		✗	✗	✗	✗	✗	✗	✗		✗	✗	✓	✓	✗	✗
Emergency pills	✗	✗		✗			✗		✗	✗	✗	✗	✗	✓	✗		✗	✗	✗	✗	✗	✗
OTHER																						
Vitamin K	✓	✓	✓	✓	✓	✓	✗		✓	✗	✓	✓	✓	✗	✗		✓	✓	✓	✗	✗	✗
Heparin	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✗	✗	✗		✓	✓	✓	✗	✗	✗
Sodium citrate	✓	✓	✓	✓	✓	✓	✓		✓	✓	✗	✓	✓	✗	✗		✓	✓	✓	✗	✗	✓
Anti Rho (D) Immune Globulin / Rhogham	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✗	✗	✗		✓	✓	✓	✗	✗	✗
Insulin – soluble	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✓	✓
Vitamin A mega dose	✓	✓	✓	✗			✓		✓	✓	✓	✓	✓	✗	✗		✓	✓	✓	✓	✗	✓
Ranitidine	✓	✓	✓	✓	✓	✓	✗		✓	✓	✓	✓	✓	✗	✓		✓	✗	✓	✓	✓	✗

Table 6.9: Pharmacy management practices by institution

	COLOMBO								GAMPAHA								KALUTHARA					
	TH CSHW	TH DMH	TH COLOMBO SOUTH	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH COLOMBO NORTH	DGH GAMPAA	DGH NEGOMBO	BH WATHUPITIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA		DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA
Availability of an indoor pharmacy/drugstore	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Accessible 24*7	✗	✗	✗	✓	✓	✓	✓		✗		✗	✗	✗	✗	✗		✓	✗	✗	✓	✗	✗
Availability of a drug inventory	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Drug inventory is updated	✓	✓	✓	✓	✓	✓	✓		✓	✓	✗	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Ordering drugs																						
Order at fixed time: annually or quarterly		✓					✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Order when drug reach "reorder level"	✓	✓	✓	✓	✓	✓	✓		✓	✓		✓										
Order when ever drugs ran out																		✓			✓	
Never order																						
PLASMA EXPANDERS	2000	1WK	200	150					A	A		1WK										
Availability of a buffer stock at pharmacy for these drugs																						
OXYTOCIN	✓	✓	✓		✓	✓	✓		✓	✗	✓	✓	✗	✗	✓		✓	✓	✓	✓	✓	✓
MGSO4	✓	✓	✓		✓	✓	✓		✓	✓	✗	✗	✗	✗	✓		✓	✓	✓	✗	✗	✗
HYDRLAZINE	✓	✓	✗		✓	✓	✓		✓	✓	✓	✗	✗	✗	✗		✓	✓	✓	✗	✓	✗
NIFIDIPINE	✓	✓	✓	✓	✓	✓	✓		✓	✗	✗	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
PLASMA EXPANDERS	✓	✓	✓				✗		✓	✓	✓	✓	✓	✓	✓		✓	✗	✗	✗	✗	✗
Availability of a qualified pharmacist 24 hours a day	✗	✗	✗	✓	✗	✗	✗		✓	✗	✗	✗	✗	✗	✗		✓	✓	✓	✗	✗	✗
Availability of a 'First-expiry-First-out' system for issuing drugs	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
The drug distribution register have columns for: Date of manufacture & date of expiry	✓	✗	✗	✗	✗	✗	✗		✓	✗	✓	✗	✗	✗	✓		✓	✗	✗	✗	✗	✗
The drugs arranged according to 'First in First out' basis	✓	✓	✗	✗	✓	✓	✓		✓	✓	✓	✗	✓	✓	✓		✓	✓	✓	✓	✓	✓
Drugs protected from moisture, heat or infestation	✓	✗	✗	✓			✓		✓	✗	✗	✗	✗	✗	✓		✓	✓	✓	✗	✗	✗
Drugs that require refrigeration stored in a functioning refrigerator	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓
Availability of a at least one functioning refrigerator	✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓	✓	✓	✓	✓

6.8 Equipment profile

This section describes the adequacy of selected equipment in relation to the number of deliveries and number of sections performed per day by institution, district and province. Number of deliveries is taken as a proxy measurement of the number of maternity patients needing care in the unit. The average number of deliveries per machine per day is presented as a measure of adequacy of resources needed for providing care.

The data gathered for a quarter in 2011 was used to examine the work load due to caesarean sections and is given in table 6.11. Data is presented only for institutions with specialised services.

The table suggests that CS packs per day are inadequate to meet the work load in many of the institutions.

Table 6.10: Average daily use of selected equipment by institution

Name of the Hospital	COLOMBO						GAMPAHA				KALUTARA			Colombo District	Gampaha District	Kalutara District	Western Province
	TH Castle Street	TH De Soya	TH Colombo South	BH Awissawella	BH Homagama	BH Mulleriyawa	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	DGH Kaluthara	BH Panadura	BH Horana				
Total number of deliveries in 2011	16180	9088	8188	4505	2437	242	9326	5483	5237	3874	8708	4162	3537	45251	24055	16716	86022
Equipments																	
Deliveries for a USS per day	8.87	4.98	5.61	3.09	6.68	0.00	3.00	7.51	14.35	10.61	11.93	5.70	9.69	24.80	10.98	9.16	14.98
Deliveries per CTG Machine per day	4.03	3.11	4.49	12.34	6.68	0.33	6.39	5.01	14.35	5.31	7.95	3.80	3.23	5.16	6.59	5.09	5.61
Deliveries for a Doppler Machine per day	14.78	3.11	5.61	6.17	6.68	0.66	12.78	15.02	14.35	10.61	11.93	11.40	0.00	6.17	13.18	15.27	11.54
Deliveries for a Labour room bed per day	1.23	0.86	1.02	1.37	0.83	0.33	1.22	1.37	1.43	1.33	2.98	1.63	1.62				
Deliveries for a delivery set per day	0.70	0.33	0.75	0.77	0.35	0.03	0.64	0.56	0.32	0.53	0.60	0.46	0.42				

Table 6.11: Average daily use of theatre facilities for obstetric care by institution

	COLOMBO						GAMPAHA				KALUTARA			COLOMBO DISTRICT	GAMPAHA DISTRICT	KALUTARA DISTRICT	WESTERN PROVINCE
	TH Castle Street	TH De Soya	TH Colombo South	BH Awissawella	BH Homagama	BH Mulleriyawa	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	DGH Kaluthara	BH Panadura	BH Horana				
LSCS	1304	365	633	494	199	11	740	411	477	332	762	357	332	3006	1960	1451	6417
Theatres	4	3	3	3	2	1	2	2	1	2	2	1	1	16	7	4	27
Theatre tables	8	7	6	9	4	1	10	6	2	6	4	2	3	35	24	9	68
LSCS kits	10	12	7	6	5	3	11	10	8	13	8	5	5	43	42	18	103
CS/operation table per day	1.81	0.58	1.17	0.61	0.55	0.12	0.82	0.76	2.65	0.61	2.12	1.98	1.23	0.95	0.91	1.79	1.05
CS/LSCS pack per day	1.4	0.3	1.0	0.9	0.4	0.0	0.7	0.5	0.7	0.3	1.1	0.8	0.7	0.8	0.5	0.90	0.69
CS/theatre per day	3.62	1.35	2.34	1.83	1.11	0.12	4.11	2.28	5.30	1.84	4.23	3.97	3.69	2.09	3.11	4.03	2.64

*dedicated theatre available

6.9 Transport and communication

Table 6.12: Transport and communication facilities by institution

Hospital	Western Province																	
	Gampaha							Colombo										
	TH RAGAMA	DGH GAMPAA	DGH NEGOMBO	DGH WATHUPITTIWALA	BH MIRGAMA	BH KIRIBATHGODA	DH MINUWANGODA	TH CSHW	TH DMH	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA	Kethumathie MH	BH Horana	BH Pimbura	DH Beruwala
Transport facilities (for 24*7 cover)																		
Total number of ambulances	5	4	4	3	1	1	1	2	3	6	4	2	3	1	1	3	NI	1
Means of repair	+	+	-	+	+	+	-	+	+	+	+	+	-	+	+	+	-	-
Funds available today for repairs	+	+	+	+	+	-	-	+	+	-	+	+	+	+	+	+	-	-
Fuel available today for transportation	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Adequate number of ambulance drivers	+	+	+	-	-	-	+	+	+	+	-	+	-	+	+	+	-	-
Absence of days when a driver was not available during the past month	+	+	+	+	-	-	+	+	+	-	-	+	+	-	+	-	-	+
Quality of the Ambulances																		
Availability of a fixed IV stand	0	4	4	3	1	1	1	2	3	1	0	0	1	1	1	0	NI	1
Fixed O2 cylinder or a mechanism to carry one	0	4	4	3	1	1	1	2	0	6	0	0	3	1	1	3	Ni	0
Source of light sufficient for resuscitation	0	0	0	3	1	1	1	2	2	6	0	0	0	1	1	3	NI	0
Ambu bag – for the ambulance	0	0	0	3	0	0	0	0	0	1	0	0	1	1	1	3	NI	0
Availability of a foot operated sucker	3	0	0	0	0	0	0	0	0	6	0	0	1	0	0	0	NI	0
Communication facilities (for 24*7 cover)																		
Availability of land phones in the facility	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Communication facility to be available 24*7	+	+	+	+	+	+	NI	+	+	+	+	+	+	+	+	+	+	+
Direct dialing facility in the LR	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

6.10. Cost Incurred by patients

Table 6.13: Costs Incurred by patients by institution

	COLOMBO																GAMPAHA										KALUTHARA										
	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH CSHW	TH DMH	TH DMH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA		TH RAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTHARA	DGH KALUTHARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA	
Cost incurred by Patients																																					
Out of Pocket expenditure- Normal delivery	1	1	1	1	1	1	1	1	1	1	1	1	1	1	0	0	1	0			1	1	1	1	1	1	1	0		0	1	1	1	0	0	1	
Drugs																																					
Investigations																																					
Other supplies																																					
A routine list of items for Normal delivery	✓	✓	✓	✓	✓	✓	✗	✗	✓	✗	✓	✓	✓	✓	✗	✗	✓	✗		✗	✓	✗	✓	✗	✗	✓	✗	✗	✗	✓	✓	✗	✗			✗	
Out of Pocket expenditure- Cesarean Section																																					
Drugs																																					
Investigations																																					
Other supplies																																					
A routine list of items for Cesarean Section																																					
Out of Pocket expenditure- Gynae. emergency																																					
Drugs																																					
Investigations																																					
Other supplies																																					
A routine list of items for Gynae. emergency																																					

No	Yes	Not relevant	Often	Rarely	Never
----	-----	--------------	-------	--------	-------

+	+	+	+	+	+
No	Yes	Not relevant	Often	Rarely	Never

Table 6.13 shows that in the overwhelming majority of institutions within the province patients have to

often bear some costs for drugs, investigations and other supplies.

Chapter 7

Human Resources for EmONC

Professionally qualified and skilled human resources are a prerequisite for the provision of high quality health care which in turn is an important determinant of health outcomes of the population. Quality performance depends on a combination of factors; availability of a sufficient number of trained personnel /professionals and their being competent, productive and responsive to patient needs.

The present assessment examined the distribution of human resources for obstetric care namely, specialist obstetricians, medical officers and midwifery staff; the term medical officer includes all MOs, SHOs and registrars but not senior registrars and intern house officers and the term midwifery staff include midwives and midwifery trained nurses. Opportunities for in-service training in relevant areas and some aspects of knowledge among selected providers were examined. The findings are presented in this chapter.

7.1 Workload

In the absence of documented staffing norms/ standards for the country for the different categories of staff working in obstetric units and labour rooms, it is difficult to comment on the work load. The exception was the number of deliveries per obstetric unit (which may be considered as a proxy for obstetrician) that is given in the Labour Room Guide (FHB 2007). However, since the distribution of human resources and work load of the different categories is an important determinant of quality of

care, the situation prevalent at the time of survey is described with a view to highlighting the variations in the work burden observed between categories of institutions, within each category as well as between districts. The findings may help in formulating guidelines and establishing norms for the different categories of institutions and staff.

The work load of a unit is dependent on many factors, such as geographic location in relation to other health institutions, easy accessibility in terms of transportation, availability of facilities as well as the duration of service availability in the particular institution and in the neighbouring institutions and perceptions of the community regarding the institution and other factors which influence personal preferences.

7.1.1: Work load of different categories of care providers

7.1.1 (a) Variation among individual hospitals in different types of institutions

The work load per institution has been calculated by dividing the number of births in the institution by the number of staff of the selected category serving in the obstetric units and labour rooms of that institution. The number of deliveries conducted under the supervision of one specialist obstetrician over a given time period would give an indication of the workload of a unit. However, in the University units where there is more than one specialist

obstetrician per unit, the proxy measure may lead to an under estimation of the workload per unit.

Tables 7.1 show the variation in work load in relation to normal deliveries in the individual institutions. The cut off values used for stratifying the workload and its' colour coding were decided upon based on the following:

According to the Labour Room Management Guide published by Family Health Bureau (2007), 300 deliveries per month is considered as the norm for an obstetric unit and this may be considered a proxy for the number of deliveries per obstetrician per month. World Health Report (2005) also gives the reference number of deliveries per year as 3600 which is equivalent to the 300 deliveries per month per unit.

The national averages for deliveries per month based on the data from the current assessment are 216 per obstetrician, 68 per MO obstetrics and 9 per midwife.

The norm described in the World Health Report, 2005, gives the number of medical officers required for the reference number of deliveries (3600 per year/300 per month) as 3 part time MOs and the number of midwives as 20. This is equivalent to 100 deliveries per MO per month and 15 deliveries per midwife. These values also have to be examined in relation to roles and responsibilities of the different categories of staff expected in the local scenario keeping in mind that WHO recommendations are a generalisation applicable to countries with a wide variation in service availability. Therefore in the present analysis 4 MOs per 300 deliveries per month, i.e. 75 deliveries per MO per month, were used taking into consideration the high workloads noted.

The cut off values used in this analysis and the data are presented so that they may serve as a basis for deriving staffing norms based on types of institution as well as work load of institutions.

Key to tables 7.1

No. of Deliveries per specialist obstetrician per month	No. of deliveries per MO-obstetrics *	No. of deliveries per midwifery staff**per month
<150	<30	<5
150-200	30-45	5 to 10
200+	45+	10+
250+	60+	15+
300+	75+	20+
No staff	No staff	No staff

** includes MOs, SHOs and registrars but not senior registrars and intern house officers*

***includes midwifery trained nurse and midwife*

Table 7.1: Work load of selected categories of care providers for normal deliveries by institution

Name of the Hospital	COLOMBO						GAMPAHA				KALUTARA			Colombo District	Gampaha District	Kalutara District	Western Province
	TH Castle Street	TH De Soysa	TH Colombo South	BH Awissawella	BH Homagama	BH Mulleriyawa	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	DGH Kaluthara	BH Panadura	BH Horana				
Total number of deliveries in 2011	16180	9088	8188	4505	2437	242	9326	5483	5237	3874	8708	4162	3537	45251	24055	16716	86022
Human resources																	
Deliveries /Specialist Obstetrician / month	224.72	151.47	170.58	187.71	101.54	20.17	129.53	228.46	218.21	161.42	362.83	173.42	147.38	171.41	167.05	232.17	190.21
Deliveries / MO-Obst**/ month	43.49	26.11	56.86	53.63	40.62	5.04	59.78	65.27	62.35	46.12	90.71	57.81	73.69	42.17	58.73	76.63	50.62
Deliveries /Midwifery staff*/ month	9.36	5.57	7.34	8.94	6.35	3.36	9.96	8.16	9.92	6.87	10.22	7.88	12.82	8.12	8.32	8.57	8.26

** MO-Obst in the specialist units

* Midwives and midwifery qualified nurses in the specialist units only

The table (7.1) highlights the unequal distribution of human resources within the province. It is seen that the workload per obstetrician and per MO per month is high in the Kalutara district as well as the DGH Kalutara compared to the other districts and other institutions. Deliveries per midwifery staff are also high in the Kalutara district.

7.2 In-service training profile

Continuing professional development of care providers is an integral component of quality assurance in health care provision and in-service training which helps them to sustain and update knowledge and skills forms an important method of achieving quality enhancement. Individual data on

7.2: Workload for caesarean sections by institution

Hospital	COLOMBO						GAMPAHA				KALUTARA			Colombo District	Gampaha District	Kalutara District	Western Province
	TH Castle Street	TH De Soysa	TH Colombo South	BH Awissawella	BH Homagama	BH Mulleriyawa	TH Colombo North	DGH Gampaha	DGH Negambo	BH Wathupitiwala	DGH Kaluthara	BH Panadura	BH Horana				
Number of sections performed for a quarter 2011	1304	365	633	494	199	11	740	411	477	332	762	357	332	3006	1960	1451	6417
Human resources for sections																	
CS / Specialist Obstetrician / month	72.44	24.33	52.75	82.33	33.17	3.67	41.11	68.50	79.50	55.33	127.00	59.50	55.33	45.55	54.44	80.61	53.48
CS / Specialist Anaesthetist / month	217.33	60.83	52.75	82.33	66.33	3.67	82.22	137.00	159.00	110.67	254.00	119.00	110.67	71.57	108.89	161.22	93.00
CS / Specialist Paediatrician/ month	217.33	60.83	105.50	82.33	33.17	3.67	123.33	68.50	79.50	55.33	127.00	59.50	55.33	100.20	59.39	69.10	76.39
CS / MO - Obstetrics / month*	14.02	4.20	17.58	23.52	13.27	0.92	18.97	19.57	22.71	15.81	31.75	19.83	27.67	11.39	19.22	26.87	15.28
CS / MO - Anaesthesia / month	28.98	11.06	6.39	18.30	8.29	3.67	6.67	11.42	14.45	11.07	14.94	9.92	11.07	13.01	9.33	12.40	11.50
CS / MO - Blood Bank / month	108.67	17.38	26.38	82.33	16.58	3.67	18.97	22.83	17.67	27.67	36.29	19.83	55.33	38.54	20.42	32.24	29.30

*Medical officers in relevant units only

7.1.2 Work load for caesarean sections

Table 7.2 shows the variation in work load for caesarean sections by district and province, by category of institution and category of staff. These rates are also partly accounted for by the wide variation in caesarean section rates between institutions and districts.

The table highlights the unequal distribution of human resources within the province. Marked differences are seen in the Kalutara district. It is seen that in Gampaha the human resource distribution is more equitable.

in-service training received during the 5 years preceding the assessment was sought from Medical Officers, Nursing Officers and Midwives working in obstetric units and labour rooms. Intern medical officers were excluded from the sample. Information was collected based on a list of programs relevant to the provision of obstetric services developed and conduct by the FHB in collaboration with the relevant professional organisations. The programs that were inquired about were: Essential New Born Care, Basic Neonatal Life Support, Advanced Neonatal Life Support, Lactation Management, Baby Friendly Hospital Initiative, Labour Room Management and Infection Control.

In addition to this, medical officers were asked about opportunities to work as an intern house officer in obstetrics, and paediatrics. They were also asked about training they may have had in blood transfusion procedures including blood grouping and matching.

The data were analysed by category of staff and by district and province and collated nationally. Details of training for each district are given in the provincial reports and this chapter presents the collated national data. In this analysis when at least 60% of staff in a given category has been trained in a particular program, coverage was considered as adequate and 40% or less was considered as inadequate.

Table 6.4 shows that over all levels of in-service training are inadequate in the whole province. In all categories of staff studied training in infection control, labour room management and the baby friendly hospital initiative has been poor.

7.2.2. In-service training profile of Medical Officers

Information on in-service training was collected from a sample of 50 medical officers randomly selected from the district. Medical officers working in obstetric wards and who is the first medical person to see patients in the obstetric ward and present at time

Table 7.3: In-service training profile of selected categories of personnel in the province

	NO-OT		NO-LR		MW-LR		NO-PNS		NO-NICU		Total	
<i>Sample</i>	388		350		329		190		300		1557	
Training	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Essential New Born Care	40	10%	180	51%	112	34%	50	26%	130	43%	512	33%
Neonatal Life Support	101	26%	200	57%	150	46%	73	38%	150	50%	674	43%
Lactation Management	61	16%	200	57%	201	61%	70	37%	175	58%	707	45%
Baby Friendly hospital Initiative	21	5%	85	24%	75	23%	36	19%	45	15%	262	17%
Labor Room Management	20	5%	115	33%	79	24%	18	9%	13	4%	245	16%
Infection Control	50	13%	45	13%	41	12%	10	5%	60	20%	206	13%

	<40% trained		40-59% trained		≥60% Trained
--	--------------	--	----------------	--	--------------

7.2.1 In-service training profile of selected categories of personnel

Table 7.3 shows the provincial in service training profile by category of staff. Training requirements may vary depending on the positions held by the different categories of staff. When at least 60% of staff has been trained in the identified program it was considered adequate and 40% and less was considered inadequate.

of data collection were given the data collection tool. Intern medical officers were excluded from the sample.

Table 7.4: In service training profile of Medical Officers

Table 7.4: In service training profile of Medical Officers

	No	%
Sample size	149	
Internship in Obstetrics	80	54%
Internship in Pediatrics	70	47%
Anaesthesia	25	17%
Blood Transfusion Procedures	36	24%
Essential New Born Care	59	40%
Neonatal Life Support - Basic	89	60%
Neonatal Life Support - Advanced	58	39%
Lactation Management	42	28%
Baby Friendly Hospital Initiative	25	17%
Labour Room Management	60	40%
Infection control	22	15%

It is seen that more than half the MOs have had training in obstetrics. Infection control, lactation management, baby friendly hospital initiative are areas where training is poor.

7.3 Knowledge among selected categories of care providers

This section present the information on knowledge of selected categories of care providers on EmONC and the analysis was done nationally.

7.3.1 Sample

Knowledge and related practices among selected categories of institutional care providers was assessed using a self-administered questionnaire. The targeted providers were Midwives, midwifery trained Nursing Officers and House Officers/ Medical officers working in the wards and the labour rooms in obstetric units, considering them as the first persons likely to come in to contact with an emergency situation in management of a woman in labour or an emergency. These categories of staff, present on duty at the time of data collection were given the questionnaires and requested to answer individually without discussion. However, since this activity had to be carried out without disturbing the functioning of the unit it was not feasible to administer the questionnaires under conditions of an academic examination. The completed questionnaires were collected by the ward sister/nurse-in-charge at the end of the shift.

7.3.2 Questionnaire

The questionnaire consisted of theoretical questions as well as questions based on a set of six case scenarios on common obstetric emergencies as well as routine management of labour namely; PPH, PIH, sudden collapse, heart disease, active management of the third stage of labour and monitoring of labour.

The instrument was designed to test different areas of knowledge and practices such as:

1. Theoretical Knowledge e.g. List causes
2. Basic steps in managing an emergency e.g. Start an infusion/ start on Magnesium Sulphate (Mg SO₄) in eclampsia
3. Tendency to be thoughtful and proactive .e.g. Informing the operating theatre and anesthetist of “possible” need for intervention in a patient with postpartum collapse before the decision to operate
4. Influence of personal attitudes in decision making in the absence of documented guide lines e.g. Informing the next level officer /informing the specialist obstetrician
5. Thinking of and responding to rare but near fatal situations e.g. inversion of uterus

7.3.3 Analysis

The data were analysed by category of staff and are presented at the national level. District and provincial analysis was not attempted because of relatively small sample sizes. Data were checked for completeness and analysed using SPSS. Each correct answer was given one mark and percentages were calculated based on the total marks that could be obtained. The percentage of marks obtained for each component of the questionnaire is also calculated using the same method. The questions on enumeration of causes are presented as the number and percentage of persons with correct responses.

7.3.4 Results

The analysis is based on a sample of 1182 care providers: 445 midwives, 439 midwifery trained nursing officers and 298 medical officers selected randomly.

7.3.4.1 Institutional midwives

Answering the question on the causes of PPH, 74% of the midwives gave 3 causes (expected no. of causes) and a further 40 % mentioned 2 causes. However with the question on PIH it was only 40 % who gave 3 correct responses with a further 19.8 % providing two correct responses. Only 40% and 38% respectively could give 3 causes for sudden collapse and symptoms of heart disease and it is noteworthy that 3-4% of midwives did not provide a single correct answer for any of the questions.

Table 7.5: The number of respondents in each category

Category of staff	No. Responded
Midwives	445
Midwifery trained nursing officers	439
Medical officers	298
Total	1182

Table 7.6: Causes given by midwives for each condition

	3 correct responses		2 Correct responses		1 correct responses		No correct answers	
N=445	No.	%	No.	%	No.	%	No	%
PPH Causes	331	74.4	88	19.8	14	3.1	12	2.7
PIH affecting Organs	182	40.9	199	44.7	50	11.2	14	3.1
Causes for sudden collapse	176	39.6	172	38.7	84	18.9	13	2.9
Symptoms of heart disease	169	38.0	204	45.8	53	11.9	19	4.3

Table 7.7: Frequency distribution of the causes of PPH as given by midwives

Cause of PPH	Number	Percentage
Retained products	187	42.0
Tears in the genital tract	161	36.2
Multiple pregnancy	149	33.5
Multi parity	146	32.8
Retained placenta	131	29.4
Anaemia	120	27.0
Uterine atonia	94	21.1
PIH	39	8.8
Placental abnormalities	33	7.4
Coagulopathy	20	4.5
Fibroids complicating pregnancy	13	2.9
Polyhydramnios	6	1.3
Abruptio placentae	6	1.3
Instrumental delivery	4	0.9
Past Section	2	0.4
Other causes	72	16.2

While 42% of the sampled midwives gave retained products as a cause of PPH, only 29%, 32.2% 33.5%, 36.2% gave retained placenta, multi parity, multiple pregnancy and tears in the genital tract as causes for PPH respectively. It is noteworthy that uterine atonia one of the common causes which needs uterine massage and oxytocics were given as a cause by only

21% and trauma/tears were mentioned by only 36%. Retained placenta and retained products were mentioned by less than half the midwives. It is important to note that less than half of the midwives have given any one of the causes. These findings point to gaps in knowledge that may negatively influence an effective and timely response in an emergency.

Table 7.8: Frequency distribution of the organs affected by severe PIH (Responses from midwives)

Organs Affected by PIH	Number	Percentage
Brain	382	85.8
Kidney	313	70.3
Liver	171	38.4
Lungs	71	16.0
Eye	40	9.0
Clotting system	1	0.2

Table 7.9: Frequency distribution of the causes for sudden collapse after delivery
(Responses from midwives)

Causes for sudden collapse after delivery	Number	Percentage
PPH	408	91.7
Uterine inversion	139	31.2
Severe PIH	120	27.0
Uterine Rupture	99	22.2
Heart diseases including MI	70	15.7
Amniotic fluid embolism	43	9.7
Vasovagal Shock	37	8.3
Anaphylaxis	21	4.7
Thromboembolism	9	2.0
Septicaemia	8	1.8
Ruptured liver	1	0.2

Table 7.10: Frequency distribution of the Symptoms of severe heart disease
(Responses from midwives)

Symptoms of severe heart disease	Number	Percentage
Shortness of breath	353	79.3
Tiredness	189	42.5
Palpitations	125	28.1
Oedema	94	21.1
Cyanosis	71	16.0
Exercise intolerance	68	15.3
Chest pain	53	11.9
Orthopnea	8	1.8
Cough	6	1.3
Haemoptysis	1	0.2

Although 79% of the respondents identified breathlessness as a symptom of heart disease it is important to note that only 1.8%, 1.3 % and 0.25 considered orthopnoea, cough and haemoptysis respectively as signs of severe heart disease. While accepting that these are uncommon presentations in routine obstetric practice, care providers need to be aware of the significance of these symptoms if they are to identify conditions such as postpartum onset of pulmonary oedema which may occur in patients with heart disease.

Some of the findings on the answers to questions on clinical scenarios are given below.

- In managing a PPH 98% mentioned that they would examine and massage the uterus and 79% would bring the emergency trolley to the bedside as a first response, a proactive step. However, on the question of informing the nursing officer immediately, only 88% answered in the affirmative. The balance 12 % who did not opt to inform is a matter for concern. Whether it indicates that there is lack of clarity in what is officially expected or is due to reluctance at individual level is not clear.
- On managing eclampsia, while 95% agreed to manage the patient in left lateral position and

50% said that they would insert a mouth gag. Surprisingly only 62% disagreed with the statement that “she be managed in a bed at the end of the ward as only that one is a bed with bars”!!

- On the question on postpartum collapse, 90% have given correct responses however the

balance 10% had agreed with the statement “give her a glass of water to drink”.

Table 7.11 shows that knowledge on labour, active management of the third stage of labour and monitoring of labour, is at a lower level relative to that of PPH and PIH. It is noted that the range of marks varied from zero to hundred.

Table 7.11: Marks obtained by the midwives for each component

Marks	PPH	PIH	Sudden collapse	Heart disease	Active Management of 3 rd stage	Monitoring of labour
Mean	79.9	77.7	82.3	81.6	53.8	66.2
Median	87.5	80.0	87.5	88.8	57.1	60.0
Mode	87.5	80.0	87.5	88.8	42.8	60.0
SD	16.6	14.6	14.8	17.2	20.5	24.8
Minimum	0.0	0.0	0.0	0.0	0.0	0.0
Maximum	100	100	100	100	100	100
Marks						
≥90.0	22.9	34.6	23.8	19.3	2.2	17.1
70.0-89.9	54.2	49.2	60.4	61.3	30.1	30.3
50.0-79.9	20.4	13.7	13.9	15.3	24.3	31.2
<50	2.5	2.5	1.8	4.0	43.4	21.3

Table 7.12: Summary of marks obtained by the midwives

Marks range	Frequency	Percentage
≥90	14	3.1
80-89	132	29.7
70-79	199	44.7
60-69	61	13.7
50-59	30	6.7
<50	9	2.0
Total	445	100

It is seen that nearly 77% of participants have scored over 70% and 98% of participants scoring over 50% marks , suggesting that overall knowledge in the areas tested are good.

7.3.4.2 Nursing officers

Table 7.13: Causes given by nursing officers for each condition

	4 correct responses		3 correct responses		2 Correct responses		1 correct responses		No correct responses	
N=439	No	%	No.	%	No.	%	No.	%	No	%
PPH Causes	313	71.3	105	23.9	15	3.4	4	0.9	2	0.5
PIH affecting Organs	86	19.6	261	59.5	69	15.7	19	4.3	4	0.9
Causes for sudden collapse	119	27.1	135	30.8	117	26.7	58	13.2	10	2.3
Symptoms of heart disease	107	24.4	185	42.1	113	25.7	24	5.5	10	2.3

In response to the question on causes of common obstetric emergencies, Nursing Officers were expected to list four causes for each condition. It is seen that, 71% of the sample were able to list 4 correct causes of PPH and a further 23 % gave 3 causes. However, only 19% gave 4 correct responses

in the case of PIH, with a further 60% giving three correct responses. On the questions on postpartum collapse and heart failure it was only 25% who gave 4 correct responses with a further 30-40% giving only 3 responses.

Table 7.14: Frequency distribution of the causes of PPH
(Responses by nursing officers)

Cause of PPH	Number	Percentage
Multi parity	226	51.5
Retained placenta	189	43.1
Multiple pregnancy	174	39.6
Tears in the genital tract	160	36.4
Anaemia	156	35.5
Retained products	140	31.9
Uterine atonia	131	29.8
PIH	94	21.4
Coagulopathy	73	16.6
Placental abnormalities	70	15.9
Fibroids complicating pregnancy	39	8.9
Polyhydroamnios	31	7.1
Abruptio placentae	14	3.2
Past Section	9	2.1
Instrumental delivery	7	1.6
Other causes	88	20.0

Of the nursing officers who responded to the questionnaire, 43%, 39%, 17% and 36% listed retained placenta, multiple pregnancy, coagulopathy and tears in the genital tract respectively as causes of PPH. Common cause of uterine atonia was mentioned only by 30%. Tears / trauma to the genital tract have not been given as a reason by the vast majority (64%). Although a majority of persons were able to give four correct responses, the omission of common conditions that need to be thought by a majority is a cause for concern.

Ninety three percent of Nursing Officers gave PPH as a cause of sudden collapse after delivery. However, uterine inversion and uterine rupture was given only by 40.8%, 28% and 13.7% of respondents respectively indicating the need to make them aware of rare but critical conditions needing immediate and specific responses.

**Table 7.15: Frequency distribution of the organs affected by severe PIH
(Responses from nursing officers)**

Organs Affected by PIH	Number	Percentage
Brain	422	96.1
Kidney	384	87.5
Liver	273	62.2
Lungs	133	30.3
Clotting system	1	0.2
Eye	61	13.9

**Table 7.16: Frequency distribution of the causes for sudden collapse after delivery
(Responses from nursing officers)**

Causes for sudden collapse after delivery	Number	Percentage
PPH	410	93.4
Uterine inversion	179	40.8
Uterine Rupture	123	28.0
Amniotic fluid embolism	120	27.3
Heart diseases including MI	99	22.6
Severe PIH	74	16.9
Vasovagal Shock	67	15.3
Anaphylaxis	60	13.7
Thromboembolism	28	6.4
Septicaemia	12	2.7

**Table 7.17: Frequency distribution of the symptoms of severe heart disease
(Responses from nursing officers)**

Symptoms of severe heart disease	Number	Percentage
Shortness of breath	371	84.5
Palpitations	199	45.3
Tiredness	195	44.4
Oedema	193	44.0
Cyanosis	106	24.1
Chest pain	84	19.1
Exercise intolerance	60	13.7
Orthopnoea	17	3.9
Cough	4	0.9
Haemoptysis	2	0.5

Table 7.17 shows that among the nursing officers only 4% considered orthopnoea as a sign they need to take note of as a symptom of severe heart disease.

Some of the findings on the answers to questions on clinical scenarios are given below.

- On the management of PPH 87.5% of the nursing officers agreed that they would call the HO/SHO immediately and take a blood sample for cross matching indicating a common and good practice which goes beyond what is considered and documented as their “duty”.
- A matter for concern is the fact that only 40% agreed that the response “if vital biological parameters are normal will wait for 15 minutes and review”, was a wrong practice.
- Seventy four percent of respondents indicated that they would inform the middle level of medical officers i.e. SHO/Registrar at the same time as the HO in case of a PPH. This indicates a good practice in an emergency that has the potential to deteriorate rapidly.
- On the question on eclampsia 97% mentioned that they would inform the HO immediately. 90% said that they would catheterize and insert a Foley catheter and start to maintain a urine output chart while awaiting the HO, a proactive and a useful step.
- In dealing with a collapsed patient nearly 75% of the nursing officers said that they would request permission from the HO to start an IV infusion at the time of informing about the emergency situation. Another proactive and good practice in an emergency.
- Nearly all answered the question on active management of labour correctly but on progress of labor there seems to be confusion with only 43% considering “good FHS as indicating good progress” as a false statement

Table 7.18 Marks obtained by the nursing officers for each component

Marks	PPH	PIH	Sudden collapse	Heart disease	Active Management of 3 rd stage	Monitoring of labour
Mean	81.4	84.4	75.9	76.3	59.6	63.9
Median	77.7	90.9	77.8	81.8	57.1	60.0
Mode	88.9	90.9	77.8	81.8	71.4	60.0
SD	14.1	12.5	17.4	15.8	21.9	24.2
Minimum	11.1	0.0	0.0	0.0	0.0	0.0
Maximum	100.00	100.0	100.0	100.0	100.0	100.0
Marks						
≥90.0	16.2	52.6	14.1	25.3	5.0	13.4
70.0-89.9	63.8	39.4	49.9	53.1	39.6	28.5
50.0-69.9	17.5	6.2	29.4	17.5	23.9	33.9
>50	2.5	1.8	6.6	4.1	31.4	24.1

It is seen that the mean as well as the modal marks for active management of the third stage of labour and monitoring of labour are lower relative to the other

areas examined. It is also noteworthy that the range of marks scored in 5 of the 6 components extends from 0-100.

Table 7.19: Summary of marks obtained by the nursing officers

Marks range	Frequency	Percentage	Cumulative percentage
≥90	16	3.6	3.6
80-89	124	28.2	31.9
70-79	202	46.0	77.9
60-69	78	17.8	95.7
50-59	8	1.8	97.5
<50	11	2.5	100.0
Total	439	100	

The summary marks (table 7.25) show that 77% of nursing officers have scored over 70%.

7.3.4.3 Medical officers

Table 7.20: Causes given by medical officers for each condition

	5 correct responses		4 correct responses		3 correct responses		2 Correct responses		1 correct responses		No responses/not responded	
N=298	No	%	No	%	No.	%	No.	%	No	%	No	%
PPH Causes	165	55.4	88	29.5	37	12.1	3	1.0	0	0.0	5	1.7
PIH affecting Organs	35	11.7	188	63.1	72	24.2	1	0.3	0	0.0	2	0.7
Causes for sudden collapse	85	28.5	106	35.6	66	22.1	25	8.4	8	2.7	8	2.7
Symptoms of heart disease	62	20.8	137	46.0	69	23.2	19	6.4	5	1.7	6	2.0

Answering the question on causes of PPH, 55% of the Medical Officers taking part in the assessment mentioned 5 causes as expected, with a further 30% mentioning four correct causes. In the other 3 areas tested viz. PIH, sudden collapse and heart disease, 5 correct responses have been given by only 12%, 29% and 21% respectively. On the question on PIH, only

12% mentioned 5 organs affected by PIH (expected) with a further 60% mentioning four correct responses. Only 30% named 5 causes (expected) of sudden postpartum collapse with only 20% naming 5 symptoms (expected) seen in heart disease in pregnancy. It is noted that 35% and 46% mentioned 4 correct responses respectively.

Table 7.21: Frequency distribution of the causes of PPH
(Responses by medical officers)

Cause of PPH (n=298)	Number	Percentage
Multi parity	211	70.8
Multiple pregnancy	175	58.7
Uterine Atonia	134	45.0
Coagulopathy	93	31.2
PIH	85	28.5
Placental abnormalities	80	26.8
Retained placenta	68	22.8
Fibroids complicating pregnancy	68	22.8
Anaemia	67	22.5
Tears in the genital tract	60	20.1
Polyhydroamnios	51	17.1
Retained products	44	14.8
Instrumental delivery	31	10.4
Past caesarean section	9	3.0
Abruptio placentae	7	2.3
Other causes	111	37.2

Looking in to the responses given on causes of PPH it is of interest to note that 55% of the medical officers did not list atonic uterus as a cause, although multi parity and multiple pregnancy two commonly documented causes of atony were given by 71% and

59% participants respectively. “Tears in genital tract (20.1%), retained placenta (22.8%) and abruption (2.3%)” were mentioned by a relatively low number of participants. Coagulopathy was mentioned by only 31%, and polyhydramnios by 17%.

Table 7.22: Frequency distribution of the organs affected by severe PIH
(Responses from medical officers)

Organs affected by PIH	Number	Percentage
Brain	297	99.7
Liver	291	97.7
Kidney	281	94.3
Lungs	188	63.1
Clotting system	19	6.4
Eye	60	20.1

Regarding the organs affected in PIH 99% mentioned brain, kidney and liver as common organs affected but only 6.4% mentioned the clotting system, a possible explanation being that they may not have considered the clotting system as an organ.

Responding to the question on symptoms of heart disease, more than 80% of medical officers mentioned breathlessness, 50% oedema but only 30% mentioned orthopnoea, a symptom indicating an advanced state of cardiac failure, which if detected and treated effectively would be lifesaving.

**Table 7.23: Frequency distribution of the causes for sudden collapse after delivery
(Responses from medical officers)**

Symptoms of severe heart disease	Number	Percentage
Shortness of breath	250	83.9
Palpitations	204	68.5
Chest pain	201	67.4
Oedema	150	50.3
Exercise intolerance	95	31.9
Orthopnoea	92	30.9
Tiredness	71	23.8
Cyanosis	28	9.4%
Haemoptysis	13	4.4
Cough	3	1.0

On causes of sudden postpartum collapse nearly 90% mentioned PPH while 50% mentioned amniotic fluid embolism but only 40% mentioned uterine rupture or inversion and only 17% considered anaphylaxis to a drug, indicating the necessity of strengthening the awareness and knowledge of these uncommon but life threatening conditions which need focused, active and immediate responses from the Medical Officers providing care during emergencies.

Some of the findings on the answers to questions on clinical scenarios are given below:

- 72% of medical officers in the sample would suspect and not give ergometrine when the scenario suggested an inversion.
- 64% of respondents would manage the basic steps of attending to a PPH and would not leave the patient to clerk a new patient.

**Table 7.24: Frequency distribution of the Symptoms of severe heart disease
(Responses from medical officers)**

Symptoms of severe heart disease	Number	Percentage
Shortness of breath	250	83.9
Palpitations	204	68.5
Chest pain	201	67.4
Oedema	150	50.3
Exercise intolerance	95	31.9
Orthopnoea	92	30.9
Tiredness	71	23.8
Cyanosis	28	9.4%
Haemoptysis	13	4.4
Cough	3	1.0

While this finding is reassuring in general, it is a cause for concern that 36% may leave the patient to attend to a routine duty.

- 54% of medical officers responding to a PPH with normal vital signs indicated that they would call the specialist obstetrician immediately. It is noteworthy that 46% did not appear to consider it as an appropriate response.
- 98% would inform the specialist obstetrician about an eclampsia patient immediately and 90% would start Magnesium Sulphate but only 46% indicated that they would start antibiotics in the presence of evidence of aspiration.
- With a post-partum collapse 95% of the participants would inform the specialist obstetrician as well as the operating theatre and anaesthetist, a proactive step and a good practice.
- In a patient with heart failure 90% would inform the specialist obstetrician and 95% would inform the ICU and only 86% would administer Frusemide. However, the finding that 5% would inform the ICU but not the specialist obstetrician raises a concern.
- More than 70% answered the questions on active management and monitoring of labour correctly.

Table 7.25: Marks obtained by the medical officers for each component

Marks	PPH	PIH	Sudden collapse	Heart disease	Active Management of 3 rd stage	Monitoring of labour
Mean	78.2	68.7	73.3	70.8	73.5	80.0
Median	80.0	72.7	80.0	72.7	71.4	80.0
Mode	80.0	72.7	80.0	72.7	85.7	100
SD	16.6	11.9	17.0	14.4	20.0	23.1
Minimum	0.0	0.0	0.0	0.0	0.0	0.0
Maximum	100.0	100.0	100.0	100.0	100.0	100.0
Marks						
≥90	36.5	5.4	26.8	9.0	15.1	42.8
70-89	46.2	48.8	48.2	56.9	59.5	31.1
50-79	14.0	39.8	18.7	28.1	12.7	16.4
>50	3.3	6.0	6.4	6.0	12.7	9.7

Table 7.31 shows that on PIH and heart disease the proportions obtaining 90% of marks or more was less than the other areas.

Table 7.26: Summary of marks obtained by the medical officers

Marks range	Frequency	Percentage	Cumulative percentage
≥90	2	0.7	0.7
80-89	64	21.5	22.1
70-79	150	50.3	72.5
60-69	61	20.5	93.0
50-59	15	5.0	98.0
<50	6	2.0	100.0
Total	298	100.0	

When considering the overall scores with nearly 72% of participants scoring over 70% and 100% of participants scoring over 50% marks the knowledge

and practices in general are satisfactory in the areas tested but it is important to fill the specific gaps described earlier.

Chapter 8

Preferences for Place of Delivery

In planning services it is important to examine women's preferences for place of delivery. It is known that for a variety of reasons women temporarily move out of their district of residence for giving birth and it is important to examine this aspect in relation to service provision. The present study collected information on the place of delivery for all births documented in their records by a midwife for a specified period of 3 months (quarter of the year in 2011). This chapter describes the preferences women have shown in choosing the place of delivery based on the field data collected together with data on births in government institutions for the year 2011.

8.1 Preferences for type of institution

Examination of published data shows that over time, births have gradually shifted from the smaller hospitals to higher level institutions. Changes in classification / grading and nomenclature of institutions make it difficult to examine these shifts by type of institution over the last decade; however it is clearly seen that there has been a gradual decline in births in institutions below the level of a base hospital. The Annual Health Bulletin 2003 notes that 18% of births in government hospitals occurred in institutions below the level of a base hospital. This proportion has gradually declined to 16% in 2005, 12% in 2007, to 5.4% in 2011.

Figure 8.1 and 8.2 are graphic presentations of data in tables 3.2 and 3.3.

Figure 8.1: Distribution of births in government institutions by type of hospital and district - 2011




Figure 8.1 shows the percentage of births in 2011 in the country by type of hospital.

Of the total 343384 births that took place in the country during 2011, nearly 40% took place in 15 TH/PGH with a further 29% taking place in 18 District General hospitals. Only 28% of births took place in the 64 base hospitals spread throughout the country. It is noteworthy that only 5.4% of births have taken place in institutions below the level of a base hospital.


In 2011, the highest proportion of births occurred in Teaching Hospitals and Provincial General Hospitals with the exception of the Ratnapura district where the births in Base Hospitals (50%) is marginally higher than in the TH/PGH type (47%) of institutions.

be related to the travel times from home to institutions.

Of the total births in government institutions only 6.9% of births occur in hospitals without specialist services. These institutions comprise of a few base hospitals, some of which have been upgraded but where a specialist obstetrician has not been appointed and hospitals below this level. The births in non specialist hospitals appear to be commoner in certain district such as Nuwara Eliya (23.6%), Moneragala (22.7%), and Mannar (21.6 %) Mullativu (19.4%), Anuradhapura (15.7%) and Batticaloa (14.4%).

Although the proportion of women delivering in institutions without specialist services is low, it amounts to 23873 of deliveries annually. This small but vulnerable group of women have the potential to

Figure 8.2: Distribution of births in government hospitals by functional classification and district - 2011



When births are examined by functional classification of institutions used in the study, it is seen that at national level 93.1 % of births occurred in hospitals where a specialist obstetrician is available. It is also noted that the preference is for larger institutions with more than one obstetric unit in districts where such facilities are available. This means that nearly 77% births occur in institutions with more than one specialist which are able to provide specialist services 24*7. The exception to this is the Trincomalee district where there is a marginally higher proportion of births in institutions with a single unit with one specialist obstetrician(49%) compared to the larger institution with more than one unit (44%). This may

generate 3581 complications per year and it is important that they have access to quality emergency obstetric services when the need arises. Therefore it is important to strengthen facilities and skills available in these institutions so that they do not become the weak link in the national chain providing EmONC.

It is noted that nearly 38730 (11.3%) deliver in hospitals where one maternity unit is shared by two specialist obstetricians. This may not be the ideal conditions for providing good quality care, leading to operational constraints and accountability issues. A further 16% (54863) deliver in institutions that have

only one specialist obstetrician. Both are situations where optimum conditions for provision of quality specialist services may not be available. Correction of these two situations through effective administrative responses will improve services to nearly 27% (93593) of women.

8.2 Movement of women between districts for delivery

It is known that recording of births by midwives may not be complete and this may be due to many reasons. Table 8.1 examines the completeness of the field data based on the births reported by the Registrar General. It is important to note that the field data was collected for a quarter in 2011 and relate to the number of women who had delivered in that quarter while the data from the RG are births registered. The analysis uses information from all Medical Officer of Health units in the country.

Although the data from the field refers to women who delivered during a quarter and the RG data to

births, it is unlikely that this factor accounts for the differences in numbers observed between the two sources. Since the percentage of twin births (0.89%) and multiple births (0.01%) (Medical Statistician, 2010) is very low, a comparison of the data from the two sources may be used to examine completeness of field data.

Field midwives may under record births taking place in the private sector and births occurring to women in urban areas. This is confirmed by the high predominance of reporting of women delivering in government institutions by the midwives (98.4%) as shown in table 8.2, This is nearly 4% more than the figure of 94.5 % reported in table 3.1 for the country based on data from the government institutions and the Registrar General. In recording births, the field midwives are also more likely to under record births that may have occurred outside their district. However, it is unlikely that women going to a particular district / districts for delivery are preferentially excluded in recording by the midwife and as such may not bias the analysis.

Table 8.1: Calculation of percentage completeness of the data reported from the field – Western Province

District	Births 2011 (RG)	Births per Quarter estimated from RG data	Deliveries in govt. hospitals	Deliveries in private hospitals	Total deliveries reported by MOH	% completeness
Colombo	54,888	13722	11136	145	11281	82.2
Gampaha	27,457	6864	5689	662	6351	92.5
Kalutara	16,947	4237	4269	158	4427	104.5
Western Province	99,292	24823	21094	965	22059	88.9
Sri Lanka	363,415	90854	80,377	1,300	81677	89.9

Table 8.2: Place of delivery reported in field data – Western Province

District	No of deliveries reported from the field	No of deliveries in Government hospitals		No of deliveries at home		No of deliveries in a Private hospital	
		No.	%	No.	%	No.	%
Colombo	8004	7859	98.2	0	0.0	145	1.8
Gampaha	8106	7439	91.8	0	0.0	667	8.2
Kalutara	4853	4692	96.7	0	0.0	161	3.3
Sri Lanka	83095	81742	98.4	20	0.02	1346	1.6

Table 8.3 is based only on deliveries reported in the field data as taking place in government institutions. The column in respect of a district shows the numbers and district of origin of persons giving birth in that district, where as a row indicates the number moving out of a district and the district in which the delivery took place. The diagonal therefore represents the numbers living in the district who have also given birth in a government institution within the same district.

It is seen that all districts have people going to adjacent districts for delivery. While in some of the districts the influx and efflux balances itself, some of the districts demonstrate a noteworthy shift from this.

Based on table 8.3, the net “migration” index was calculated for each district. **Net migration** rate in this instance is the difference between people who come in to a district to deliver and those who go out of the district, during a given period of time presented as per 1000 deliveries reported to have occurred in the district. A positive value represents more people entering the area than leaving it, while a negative value means that more people are leaving than entering it. This is an indication of the net effect of “migration” of mothers to and from a district to the total reported deliveries of the district.

$$\text{NMI} = \frac{(\text{mothers going out of the district to deliver}) - (\text{mothers from other districts coming in to deliver})}{\text{Number of mothers reported to have delivered in the district}}$$

The NMI for districts are presented in table 8.4.

Table 8.2: Place of delivery reported in field data – Western Province

	Colombo	Gampaha	Kalutara	Kandy	Matale	Nuwara Eliya	Galle	Matara	Hambanthota	Jaffna	Killinochchi	Mannar	Vavuniya	Mullativu	Batticaloa	Ampara	Trincomalee	Kurunegala	Puttlam	Anuradhapura	Polonnaruwa	Badulla	Monaragala	Rathnapura	Kegalle
Colombo	7785		72					1													1				
Gampaha	1960	5390	2	8	2	3	2	1										46		6	1	1		3	14
Kalutara	550	2	4052	1		2	65	2	1				1			1		1		3		3	1	5	2
Kandy	12	9	3	4714	71	5	3	3	1	1			2			3	1	40	1	6	12	172	4	1	18
Matale	13	5	1	77	1710	3	2	1	1				3			2	1	70		16	9	150	4	6	5
Nuwara eliya	6	10	1	803	2	2817	2	4		1								3		2	2	23	1	1	
Galle	14	2	67	1			4231	34	5									1				3	1	3	2
Matara	10	5	4		1		294	2709	52									59		2	1	1	3	49	3
Hambanthota	11						75	459	1889									4		3			21	147	
Jaffna	1									2527	3	3	3	2								1		1	
killinochchi										66	461		3												
Mannar												351	24							9					
Vavuniya	1			1						6		1	639							2					
Mullativu	1									64	178	1	78	270	1										
Batticaloa															2201	30									
Ampara	4			12			2	3	1							3169		3		1	16	36	38	4	2
Trincomalee	3	1		16		1	2				1		1		4	2	1748	6		49	2				1
Kurunegala	28	55	1	32	81	1	5	2	1							1	1	5303	387	305	8	3	2	8	49
Puttalam	17	188	2	1	1					1					1			112	2470	7					4
Anuradhapura	2			1	230			1					101				46			3761	17				
Polonnaruwa	3			1	192			3								32	3			5	2305			1	
Badulla	4	1		7	2	91	2	2	5						1	4		2		1		3000	14	9	2
Monaragala	5								188							67						172	1558	196	
Rathnapura	294	1	64	3	1	3	2	12	6				1			1		3		2	1	4	4	3771	2
Kegalle	420	20		271				2																71	2617

* a column shows the numbers and district of origin of persons giving birth in that district
a row indicates the number moving out of a district and the district in which the delivery took place

Table 8. 4: Deliveries occurring within the district, going out and coming in to the district- Western Province

District	Deliveries in government hospitals	Delivering within the district		Going outside the district to deliver		Coming in to the district to deliver		Net migration per delivery	Net “migration” Index – NMI*per 1000 deliveries
		No.	%	No.	%	No.	%		
Ratnapura	4200	3771	89.8	404	9.6	505	12.0	0.02	20
Kegalle	3405	2617	76.9	784	23.0	104	3.1	-0.20	-200
Sri Lanka	83095	71448	86.0	10294	12.4	10294	12.4	0.00	0

*Net “migration” Index – NMI

The present survey did not examine the reasons for movement away from the district of usual residence. It may be for a variety of reasons; personal as well as related to perceptions regarding services. It may be that in some areas the closest institution or the institution most accessible in terms of public transportation is across a district boundary. Spatial information necessary for such an analysis was unavailable.

The figure 8.3 maps the movement of women between districts. The proportions of women living in the district who have delivered in a government institution within the same district are given in black. The figures in red indicate the movement depicted by the arrow.

92



Chapter 9

Family Planning Services in Government Hospitals

In the present survey a special module was developed to examine functioning of the existing institution based family planning services. Data were collected with the intention of identifying ways of improving the services. Tables 9.1-9.3 present information on the different aspects of services and are mostly self-explanatory; however a few important findings are discussed in the text.

It is noted that three institutions in the province namely, the De Soysa Maternity Home, DGH Negambo and DGH Kalutara do not have family planning clinics. Three institutions in the province have daily clinics and all others weekly clinics. A range of contraceptive methods are available except at the BH Horana where common commodities such as DMPA and IUDs are not available. Emergency contraceptive pills are available only at the Castle Street Hospital for Women.

Table 9.1 Institution based family planning services in Western province

INSTITUTION	COLOMBO								GAMPAHA								KALUTARA					
	TH CSHW	TH DMH	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA	TH RAGAMA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTUWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTARA	BH PANADURA	BH HORANA	BH PIMBURA	DH MATHUGAMA	DH BERUWALA		
FAMILY PLANNING CLINIC																						
Availability of a regularly functioning FP clinic	✓	✗	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓		✓	✗	✓	✓	✓	✓	✓	✓	✓
Frequency of the FP clinic (weekly)	DAILY		DAILY	wkly	DAILY	Wkly	Wkly	Wkly	Wkly	Wkly	Wkly	Wkly		Wkly		Wkly	Wkly	Wkly	Wkly	Wkly	2wkly	
Person who conducts the FP clinic	MO		MO	VOG	NO	NO	MO	MO	VOG/ MO		MO	MO		MO		MO	MO	MO	MO/N O	MO		
The contraceptives methods provided at the FP clinic																						
Pills	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓		✗		✓	✓	✓	✓	✓	✓	✓
Condoms	✓		✓	✓	✓	✓	✓	✓	✓		✗	✗		✗		✓	✓	✓	✓	✓	✓	✓
DMPA	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓		✓		✓	✗	✓	✓	✓	✓	✓
IUD	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓		✓		✓	✗	✓	✓	✓	✓	✓
Implants	✓		✓	✓	✗	✗	✗	✗	✗		✗	✗		✗		✓	✗	✗	✓	✓	✗	✗
ECP	✓		✗	✗	✗	✗	✗	✗	✗		✗	✗		✗		✗	✗	✗	✗	✗	✗	✗
Person who inserts IUDs at the clinic	MO/NO		MO	MO	NO	MO/NO	MO	MO	MO		MO	MO		MO		MO	MO	MO	MO	MO	NO	
Whether had training on IUD insertion	✓		✓	✓	NO	✓	✓	✓	✗		✗	✓		✓		✓	✓	✗	✓	✓	✓	✓
Person who insert the implants at the clinic	MO		MO	MO												MO			MO			
Whether had training on Implant insertion	✓		✓	✓												✓			✓			
No. referred from the ward to the FP clinic during the last 2 months	800		DNA	DNA	24	17	4	0	10		0	DNA		DNA		DNA	0	0	3	DNA		
Adequacy of privacy for patients undergoing IUD insertion	✓		✓	✓	✓	✓	✓	✓	✓		✓	✓		✓		✓	✓	✓	✓	✓	✓	✓

Table 9.2 Institution based family planning services in Western province

INSTITUTION	COLOMBO							GAMPAHA							KALUTARA					
	TH CSHW	TH DMH	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA	TH RAGAMA	DGH GAMPAAHA	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTARA	BH PANADURA	BH HORANA	BH PTMBURA	DH MATHUGAMA	DH BERUWALA
FAMILY PLANNING SERVICES																				
Provision of <u>FP counseling</u> for post partum and post abortion patients before discharge from the ward	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
Provision of <u>FP services</u> to post partum and post abortion patients before discharge from the ward	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
If not, place referral for the FP services																				
Hospital FP clinic	✓	✓	✓	✗	✓	✗	✓	✗	✓	✗	✗	✓	✗	✗	✗	✗	✓	✓	✓	✗
FP clinic at the MOH	✓	✓	✓	✓	✓	✗	✓	✓	✗	✗	✗	✗	✗	✗	✗	✓	✗	✓	✗	✓
Field PHM	✓	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✓	✓	✓	✓
Pharmacy	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗
General practitioner	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗
Availability of dedicated theatre time for permanent FP methods (Sterilization)	✗	✗	✓	✓	✓	✗	✗	✓	✗	✗	✗	✗		✗	✗	✓	✗	✗	✗	✗
If not, Arrangement for sterilization (LRT) of women																				
Routine list	✓	✓				✓			✓	✓	✓			✓		✓				
Casualty list	✓	✓							✓	✓										
Appointment (waiting list)	✓	✓																		
Availability of a trained doctor who can perform ligation and resection of tubes (LRT)	✗	✗	✓	✓	✓	✓	✗	✓	✓	✓	✓	✗	✗	✗	✓	✓	✓	✗	✗	✗
Availability of a trained doctor who can perform vasectomy	✗	✗	✗	✓	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✓	✓	✓	✗	✗	✗
No. of post partum LRT was done in your facility during the last 6 months	67	16	DNA	152	25	19		DNA	202	DNA	49	DNA	DNA		DNA	180	DNA			
% of patents under went LRT after CS	14	15.2	13.3	10	25	0		20	5	15	10				15	10	0			

Although a majority of institutions have indicated that they provide counselling services to post-partum and post abortion women only some institutions

have indicated that they provide services. Referral to the hospital family planning clinic or to field services is also poor.

Table 9.3: Logistics and documentation of the Family Planning services

INSTITUTION	COLOMBO							GAMPAHA							KALUTARA					
	TH CSHW	TH DMH	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	BH MULLERIYAWA	DH PILIYANDALA	TH RAGAMA	DGH GAMPAA	DGH NEGOMBO	BH WATHUPITTIWALA	BH MIRIGAMA	BH KIRIBATHGODA	DH MINUWANGODA	DGH KALUTARA	BH PANADURA	BH HORANA	BH PTMBURA	DH MATHUGAMA	DH BERUWALA
LOGISTICS AND DOCUMENTATION																				
NO shortages of contraceptives during the last 6 months	0		1	0	0	0	1	0	1		0	0	0	0		1	0	0	0	0
What were the shortages?																				
Pills																				
Condoms			✓				✓													
DMPA																				
IUD									✓											
Implants			✓													✓				
ECP																				
From where, the contraceptive supplies obtained																				
MOH																				✓
RMSD				✓	✓	✓	✓	✓	✓		✓	✓	✓	✓		✓	✓	✓	✓	
FHB	1		✓		✓						✓	✓	✓	✓						
Other																				
Availability of the monthly contraceptive stock return/request (H 1158)	✓		✗	✓	✗	✓	✓	✓	✓		✗	✓		✓		✓		✗		✓
Regularly send the monthly return (H 1200 A) to MOH				✓	✓	✗	✓	✓	✓		✓	✓		✓		✓		✗	✓	✓
Any methods of following up the clients	✓		✓	✓	✓	✗	✗	✗	✗		✓	✓		✓		✓		✓	✓	✓

Chapter 10

Neonatal Care Services

Neonatal care services in the country are in the process of being organised. The information collected has been planned with a view to providing inputs in to this process.

10.1 Distribution of neonatal care facilities within the province

Seven Neonatal Intensive Care Unit (NICU) are available in the Western Province. All the institutions

that have NICU facilities have built in Special Care Baby Units (SCBU). Data for Sri Jayawardenapura Teaching Hospital was not available. BHs Mulleriyawa, Kiribathgoda, Meerigama, Pimbura and Matugama are recently upgraded Base Hospitals and do not have separate Special Care Baby Unit Facilities. Mother Baby Centres (MBC) and Lactation Management Centres (LMC) should be available in all the specialist hospitals.

Table 10. 1a Facilities available for neonatal care in the Western Province

	CSHW	DMHW	LRH	SJGH	Colombo South TH	BH Avissawella	BH Homagama	BH Mulleriyawa
NICU	1	1	1		1			
SCBU	1	3	1		1	1	1	0
Incubators at NICU and SCBU/SCBU	13	22	10		8	6	6	
Ventilators at NICU & SCBU / SCBU	4	4	5		4	1	1	
CPAP at NICU & SCBU / SCBU	3	2	1		2	0	2	
Cots in the SCBU	55	16	15		7	15	9	
Rest room for mothers of babies in NICU & SCBU / SCBU			✓				✓	
Mother baby centre								
MBC	1	0	0		0	0	0	0
Number of beds	5							
Lactation Management Centre								
LMC	1	1	1		1	1	0	0
Dedicated staff for LMC			✓		✓	✓		

Table 10. 1b Facilities available for neonatal care in the Western Province

	TH Ragama	DGH Gampaha	DGH Negombo	BH Wathupitiwela	BH Meerigma	BH Kiribathgoda	DGH Kalutara	BH Panadura	BH Horana	BH Pimbura	BH Mathugama
NICU	2	1					1				
SCBU	2	1	1	1	0	0	1	1	1	0	0
Incubators at NICU and SCBU/SCBU	13	5	5	6			14	11	8		
Ventilators at NICU & SCBU / SCBU	6	1	1	0			3	2	0		
CPAP at NICU & SCBU / SCBU	0	0	0	0			2	0	0		
Cots in the SCBU	17	8	15	5			10	12	8		
Rest room for mothers of babies in NICU & SCBU / SCBU			✓	✓			✓	x			
Mother baby centre											
MBC	0	1	1	1	0	0	1	0	1	0	0
Number of beds		3	12	8			14		6		
Lactation Management Centre											
LMC	1	0	1	1	0	0	1	1	1	0	0
Dedicated staff for LMC	✓		✓	✓			✓		✓		

Available
 Not Available
 DNA
 Data Not Available

Although no indicators have been published for neonatal care availability and geographic distribution of neonatal care services were examined in the same manner as described for EmOC.

10.2 Availability and geographic distribution of neonatal care facilities

Availability of services was examined in terms of population bases on the standard given for Emergency Obstetric and Neonatal Care, the premise being that there should be at least five neonatal care facilities for a population of 500,000 of which at least one should be a NICU and the others SCBUs.

It is important that facilities for sick neonates should be available on a 24*7 basis. Therefore the ability of institutions with NICU and SCBU facilities to provide these services on a 24*7 basis were assessed based on a set of criteria agreed upon through a consultative process and approved by the steering committee.

Also average land area per facility ad provides some indication about access within a district/province. This is a crude indicator of access in that it does not provide any information on the spread of institutions within the district.

All three of the above indicators are given in table 10.2.

Table 10.2: Availability of NICU / SCBU by population and extent

	Colombo	Gampaha	Kalutara	Western Province
Total enumerated population – 2012	2,323,826	2,298,588	1,214,880	5,837,294
Total habitable land area Km ²	699	1387	1598	3,684
Facilities: NICU	4	3	1	8
Facilities: SCBU*	6	5	3	16
NICU / 500000 population	0.9	0.7	0.4	0.7
Area in km ² /NICU	174	442	1598	460.5
SCBU / 500000 population	1.3	1.1	1.2	1.2
Area in km ² /SCBU	1/87.3 km ²	1/277.4km ²	1/532.6 km ²	1/230.2 km ²
Institutions; Specialist Neonatologist available	2	0	0	2
Institutions: Specialist Paediatrician holding position of Neonatologist available	4	2	2	8
Institutions: 2 or more Specialist Paediatrician/Neonatologist available	6	4	3	13
No. of Specialist Paediatricians** + Neonatologists available to provide neonatal care	16	10	6	32
No of Institutions with Specialist Paediatricians	7	4	3	14
Institutions with a Specialist Paediatrician / 500,000 population	1.5	0.87	0.82	1.19
Institutions with a Specialist Paediatrician / area	1/99.8 km ²	1/335 km ²	1/525.3 km ²	1/263 km ²
Facilities: 24*7 NICU	4	3	1	8
Facilities: 24*7 SCBU	5	5	3	13
24*7 NICU / 500000 population	0.9	0.7	0.4	0.7
24*7 NICU / km ² area	174	442	1598	460.5
24*7 SCBU / 500000 population	1.3	1.1	1.2	1.2
24*7 SCBU /km ² area	116.5	277.4	532.7	263.1

*All the institutions with NICU also have SCBUs. These SCBUs are calculated as separate units in this calculation.

** Only the Specialist Paediatricians holding position of Neonatologist, Specialist Paediatricians covering up for Neonatologist and in hospitals without Neonatologist the Specialist Paediatricians providing neonatal care are included in this calculation. BH Mulleriyawa has a Specialist Paediatrician.

24*7 criteria for NICU

1. Functional / operational NICU / SCBU
2. Electricity
3. Functional generator
4. Continuous source of water
5. At least two Specialist Pediatricians
6. At least six MO NICU/SCBU

24*7 criteria for SCBU

1. Functional / operational SCBU
2. Electricity
3. Functional generator
4. Continuous source of water
5. At least two Specialist Pediatricians
6. At least three MO SCBU

Two Specialist Neonatologists are available within the Western Province. Specialist Paediatrician assigned to the post of Neonatologist are available in all the districts of the Western Province. In the absence of the Neonatologist one of the other Specialist Paediatricians cover up duties at the NICU and SCBU.

10.3 Morbidity profile

Information on the morbidities seen in the neonatal units were obtained through a prospective survey carried out for one month. Tables 10.3 present the data obtained from the NICUs and SCBUs.


Whenever there is a NICU a SCBU is incorporated into it. Therefore the morbidity data from institutions with NICUs also includes the admissions needing SCBU care.

The figure 10.1 show the pattern of transfer of newborns from the neonatal units of the specialist institutions in the Southern Province. It is noted that all the transfers during the study period followed the institutional hierarchy within the district. Newborns requiring highly specialised care are transferred to the TH Kandy.

Table 10.3 Morbidity pattern at the NICU and SCBU in the Western Province

Reason for admission	Admissions to NICU	Admissions to SCBU
No of tickets reviewed	471	233
Birth Asphyxia	42	04
Meconium aspiration	28	05
Septicaemia	67	53
Hypoglycaemia	54	22
Jaundice	120	48
Idiopathic Respiratory Distress Syndrome (IRDS)	99	08
Congenital Pneumonia	12	02
Meningitis	09	05
Persistent Pulmonary Hypertension (PPHN)	05	00
Intraventricular Haemorrhage/Periventricular Haemorrhage (IVH/PVH)	01	00
Necrotising Enterocolitis (NEC)	06	03
Other	235	95

Figure 10.1 Pattern of neonatal transfers in the Western Province



The figure 10.1 show the pattern of transfer of newborns from the neonatal units of the specialist institutions in the Southern Province. It is noted that all the transfers during the study period followed the institutional hierarchy within the district. Newborns requiring highly specialised care are transferred to the Lady Ridgeway Hospital.

10.4 Interventions performed at the NICUs and SCBUs in the Western Province

Tables 10.4 provide information on the type of interventions carried out in the different level of institutions. This information was based on the 4 week prospective data collection.

Table 10.4 Interventions performed at the NICUs and SCBUs in Western Province

Intervention	Interventions performed in the NICUs	Interventions performed in the SCBUs
No of tickets reviewed	471	233
Ventilation	42	07
Nasal Continuous Positive Airway Pressure Ventilation (CPAP)	45	00
Cardio-pulmonary resuscitation	45	07
Use of phototherapy in the first 24 hours	19	24
Exchange transfusion	06	01
Any surgery	03	00
Surfactant administration	20	03
Use of any inotropes	63	00
Use of anticonvulsants	32	05
Use of therapeutic intravenous antibiotics	299	84
Use of steroids to treat refractory hypoglycaemia	03	01
Use of Glycogen	04	00
Use of 12.5% Dextrose	44	08
Use of any blood products	78	07
Use of Umbilical Artery Catheter (UAC), Umbilical Venous Catheter (UVC), Arterial/Long lines	70	03
Other	34	44

The table 10.4 shows that ventilation, cardiopulmonary resuscitation, the use of surfactant, inotropes, anticonvulsants and blood products are more in the NICUs and these reflects the morbidity as well as facilities available in these.

10.5 Resources for neonatal services

10.5.1 Infrastructure

Standard guidelines for the establishment of infrastructure facilities are given in the Building and other guidelines for NICU, SCBU, MBC (FHB 2007). Availability of essential areas as per the guidelines is indicated in table 10.5.

None of the NICUs and SCBUs had all the areas as stipulated in the type plan as given in the National Guidelines for NICUs and SCBUs (Family Health Bureau 2007). A separate room for mothers of babies in the NICU or SCBU was available only in five units.

10.5.2 Human resources

Neonatal Care is provided by specialist Neonatologist or specialist Paediatricians, Medical Officers in the NICU/SCBU and the nursing Officers in the NICU/SCBU. Still there are no Specialist Neonatologists appointed to the Western Province and at the time of the survey a Specialist Paediatrician was appointed to the cadre position of a Specialist Neonatologist. In the hospitals without cadre position for Neonatologist the Specialist Paediatrician in the hospitals provide services for the neonates as well.

With regard to Medical Officers most of the hospitals have designated Medical Officers for the NICU/SCBUs. In the hospitals without designated officers, the medical officers attached to the Paediatric Wards carry out the functions of the MO NICU/SCBU.

Table 10.5: Infrastructure for neonatal services by institution and district

	COLOMBO										GAMPAHA				KALUTARA		
	TH CSHW	LRH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAAH	DGH NEGOMBO	BH WATHUPITIWALA	DGH KALUTARA	BH PANADURA	BH HORANA
UFI	L101		L102	L102	L102	L104	L104	L105	L106	B101	B101	B302	B203	B304			
ARRANGEMENT OF THE NICU/SCBU																	
Porch	✓	✗	✗	✗	✗	✗	✓				✓	✓	✗		✗	✗	✗
Trolley transfer & reception area	✓	✓	✗	✗	✗	✓	✗		✗	✓	✓	✓	✗	✓	✓	✓	✓
Scrubbing area	✗	✗	✓	✗	✗	✗	✓		✗	✓	✓	✓	✗	✗	✓	✗	✗
Stabilization area	✗	✓	✗	✗	✗	✓	✗		✗	✓	✓		✗	✗	✓	✗	✗
High Dependency area	✓	✓	✗	✗	✗	✗	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✗
Low dependency area	✓	✓	✗	✗	✗	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✗
Procedure room	✗	✗	✗	✗	✗	✗	✗	✗	✓	✓	✗	✗	✗	✗	✗	✗	✗
Duty station	✓	✓	✗	✗	✗	✓	✗		✓	✓	✓	✗	✓		✓	✓	✓
Isolation section	✓	✗	✗	✗	✗	✓	✗	✓	✗	✗	✗	✓	✗	✓	✓	✓	✓
Incubator washing and drying area	✓	✗	✗	✗	✗	✓	✗	✓	✗	✗	✗	✓	✗	✓	✓	✓	✓
Sterilization room	✓	✓	✗	✗	✗	✗	✗	✗	✓	✗	✓	✗	✗	✓	✓	✓	✓
Sterilized good stores	✓	✓	✗	✗	✗	✗	✗	✗	✓	✓	✓	✓	✗	✓	✓	✓	✓
Pantry	✗	✓	✓	✗	✗	✗	✗	✗	✗	✓	✓	✓	✗	✓	✓	✗	✗
Drug store room	✓	✓	✓	✓	✗	✗	✗	✓		✓	✓	✗	✗	✓	✓	✗	✗
Linen washing room	✗	✓	✗	✗	✗	✗	✗		✓	✗	✗	✗	✗	✗	✓	✓	✓
Cleaners room	✓	✓	✗	✗	✗	✓	✗		✗	✗	✗	✗	✗	✗	✓	✓	✓
Dirty utility room	✓	✗	✗	✗	✗	✗	✗		✗	✗	✓	✓	✗	✓	✓	✓	✓
Veranda	✓	✓	✗	✗	✗	✗	✗		✓	✓	✓	✓	✓	✓	✓	✓	✓
Mother baby unit	✓	✗	✗	✗	✗	✗	✗	✓	✗	✗	✗	✓	✓	✓	✗	✗	✗
Breast feeding room	✗	✓	✗	✓	✗	✗	✗	✓	✗	✗	✗	✗	✓	✗	✓	✓	✓
Counseling room	✗	✓	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗
Toilets for MBU	✓	✗	✗	✓	✗	✗	✗		✓	✗	✗	✓	✓	✓	✓	✓	✓
Dining room for MBU	✓	✗	✗		✗	✗	✗		✗	✗	✗	✗	✓	✓	✓	✓	✓
Consultants room	✓	✓	✗	✗	✗	✗	✗	✗	✓	✗	✗	✓	✓	✗	✓	✓	✓
MO/NICU room	✓	✓	✓	✗	✗	✗	✗	✗	✗	✗	✓	✓	✓	✓	✓	✓	✓
Sister's room	✓	✓	✓	✗	✗	✗	✗	✓	✗	✗	✓	✓	✓	✗	✓	✓	✓
Nurses room	✓	✓	✓	✓	✗	✗	✗	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓
Minor staff room	✓	✓	✗	✗	✗	✗	✗	✓	✗	✓	✓	✗	✓	✓	✓	✓	✓
Store room	✓	✗	✗	✗	✓	✓	✓	✓	✓	✗	✗	✓	✓	✓	✓	✓	✓
Gas room and unloading bay	✗	✗	✗	✗	✗	✓	✗	✓	✓	✗	✗	✗	✓	✗	✗	✓	✗

Table 10.6 Human resource distribution by district and province

	Colombo District	Gampaha District	Kalutara District	Western Province
Total reported deliveries in Government hospitals	45251	24055	16716	86022
No of babies needing Special Care*	6787	3608	2507	12903
No. of Specialist Neonatologists available	2	0	0	2
No. of Specialist Neonatologists and Paediatricians caring for neonates**	16	10	6	32
No. MO/NICU and /or SCBU	69	37	19	125
No. of nurses at NICU and / or SCBU	194	80	56	330
Births per Neonatologist/Paediatrician	2828	2406	2786	2688
NICU &/or SCBU admission per Neonatologist/Paediatrician per month	43	34	35	39
Births per MO/NICU and /or SCBU	656	650	880	688
NICU &/or SCBU admissions per MO/NICU and /or SCBU per month	10	09	11	10
NICU &/or SCBU admissions per nurse at NICU and /or SCBU per month	4	4	4	4

*15% of the deliveries are likely to need special care (Building and Other Guidelines for Neonatal Intensive Care Units, Special Care Baby Units and Mother Baby Centers, Family Health Bureau, 2012)

**The Specialist Neonatologists, Specialist Paediatricians holding position of Neonatologist, Specialist Paediatricians covering up for Neonatologist and in hospitals without Neonatologist the Specialist Paediatricians providing neonatal care are included in this calculation

10.5.3 Guidelines, protocols, records and registers

Tables 10.7 and 10.8 indicate the availability of guidelines and protocols for the management of common newborn morbidities and the routine registers that should be available in the NICUs and SCBUs.

It is seen that there is a wide variation in the availability of guidelines and protocols in the neonatal units in the Western Province.

Table 10.7: Availability of protocols and guidelines at the NICUs and SCBUs

	COLOMBO									GAMPAHA					KALUTARA		
	TH CSHW	LRH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITIWALA	DGH KALUTARA	BH PANADURA	BH HORANA
UFI	L101		L102	L102	L102	L104	L104	L105	L106	B101	B101	B302	B203	B304			
MANAGEMENT PROTOCOLS FOR																	
Hypoglycaemia	✓	✗	✗	✓	✓	✓	✗	✓	✓	✓	✓	✓	✗	✗	✓	✗	✗
Birth asphyxia	✓	✗	✗	✗	✓	✓	✗	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓
Septic baby	✓	✗	✗	✗	✗	✓	✗	✓	✓	✓	✓	✗	✗	✗	✗	✗	✗
Meconium Aspiration Syndrome	✓	✗	✗	✗	✓	✓	✗	✓	✓	✓	✓	✗	✗	✗	✗	✗	✗
Persistent pulmonary HT	✓	✗	✗	✗	✗	✓	✗	✓	✓	✓	✓	✗	✗	✗	✗	✗	✗
IRDS	✓	✗	✗	✗	✓	✓	✗	✓	✓	✓	✓	✗	✗	✗	✗	✗	✗

Table 10.8: Registers and records available at NICU / SCBU

	COLOMBO									GAMPAHA					KALUTARA		
	TH CSHW	LRH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITIWALA	DGH KALUTARA	BH PANADURA	BH HORANA
UFI	L101		L102	L102	L102	L104	L104	L105	L106	B101	B101	B302	B203	B304			
REGISTERS AND RECORDS																	
NICU/SCBU register	✓	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Neonatal transfer form	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Neonatal diagnosis card	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Neonatal monthly return	✓	✗	✗	✗	✓	✓	✓	✓		✓	✓	✓	✗	✗	✓	✗	✗
Discharge register	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Death register	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Statistics file	✓	✓	✗	✓	✗	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓
Correspondence Book/ File	✓	✓	✗	✗	✗	✓	✓	✓	✓	✓	✓		✗	✗		✗	✗
Perinatal conference minutes file	✗	✓	✗	✗	✗	✗	✓	✓	✓	✓	✗	✗	✗	✓	✗	✗	✓
Neonatal Drug Doses Book	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

Availability of Newborn Formats that were introduced to the system in 2008 is satisfactory in more than 50% of the hospitals. The minutes of the Perinatal Conference had been maintained in seven out of seventeen hospitals in the province.

10.5.4. Selected practices related to neonatal care Neonatal Examination

Neonatal examination format (H 1162) is the Bed Head Ticket of the newborn. At birth irrespective of the place of birth (labour room, open theatre) a newborn examination format is initiated and is attached to the mother's BHT.

It is mandatory that all the newborns are examined by a Medical Officer prior to discharge from the hospital and recorded in the Neonatal Examination Format and the Child Health Development Record.

Mandatory Newborn Examination before discharge is performed in all the institutions in the Western Province and makes the necessary referrals. The Neonatal Examination Format where the examination findings are recorded was available in all the hospitals in the province. Clinic follow up plans are also available for the newborns discharged from all of the institutions in the Western Province.

Readiness for neonatal resuscitation

The neonatal resuscitation trolley, its availability and its contents were used as a proxy for measuring of readiness for neonatal resuscitation. This was assessed for each institution and is summarized and presented by type of hospital.

Table 10.9: Selected practices related to neonatal care by institution and district

	Colombo							Gampaha				Kalutara		
	CSHW	DMHW	LRH	SJGH	TH Kalubowila	BH Avissawella	BH Homagama	TH Ragama	DGH Gampaha	DGH Negombo	BH Wathupitiwela	DGH Kalutara	BH Panadura	BH Horana
UFI														
Record keeping														
Use the neonatal examination format	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Use NICU/SCBU history record sheet	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
The information recorded in the mothers BHT	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Neonatal follow up														
Neonatal examination by MO before discharge	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Referrals for the necessary conditions (eye, ENT)	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Clinic follow up plan	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Use neonatal transfer forms	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Practices in transferring newborns														
Ensure service availability at receiving end	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Inform the receiving end	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Accompanied by a responsible person	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Cost incurred by patients														
Out of pocket expenditure for NICU/SCBU care														
Drugs														
Investigations														
Other supplies														
Routine list for NICU/SCBU admission			0	0			0	0	0		0	0	0	0
No out of pocket expenditure for an emergency	✓	✗	✓	✗	✗		✓	✓	✗		✓	✓	✗	✗
	No out of pocket expenditure							Often						
	Has to incur out of pocket expenditure							Rarely						

Table 10.10 Drugs and equipment available for neonatal resuscitation at NICUs and SCBUs

	COLOMBO									GAMPAHA					KALUTARA		
	TH CSHW	LRH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTIWALA	DGH KALUTARA	BH PANADURA	BH HORANA
UFI	L101		L102	L102	L102	L104	L104	L105	L106	B101	B101	B302	B203	B304			
NEONATAL RESUSCITATION TROLLY																	
Neonatal Ambu (ventilatory) bag (250ml) with reservoir and O2 masks	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Infant laryngoscope with spare bulb & batteries/Straight blade	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Endotracheal tubes- 2.5, 3.0, 3.5 with introducer stilette	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Straight suction catheter with thumb port-FG 5,6,8,10	✓	✓	✗	✓	✗	✗	✗	✓	✓	✓	✓	✗	✓	✗	✗	✓	✗
Feeding tube – 5,6,7,8,	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Emergency tray with																	
syringes 1,2,5,10 ml	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓
Cannula 23,25 G	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Umbilical catheterization pack	✓	✓	✗	✓	✗	✓	✗	✓	✗	✗	✓	✗	✗	✗	✗	✗	✗
Blood sample bottles, culture bottles	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓
Drugs																	
Adrenaline	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Nalaxone	✓	✓	✓	✓	✗	✓	✓	✓	✓	✗	✓	✓	✗	✓	✓	✗	✓
N.Saline	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
8.4% Na HCO ₃	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
10% dextrose	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✗	✗	✓	✗	✗	✓
Distilled water	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

10.5.5 Drugs and equipment

Availability of essential drugs and equipment for neonatal care are given in Tables 10.11 – 10.13.

All SCBUs and NICUs should have infrastructure facilities for infection control. The availability of these facilities by the institution and district are presented in Table 10.12.

Table 10.11 Availability of essential drugs and equipment by institution and district

	COLOMBO									GAMPAHA					KALUTARA		
	TH CSHW	LRH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTIWALA	DGH KALUTARA	BH PANADURA	BH HORANA
UFI	L101		L102	L102	L102	L104	L104	L105	L106	B101	B101	B302	B203	B304			
Drugs available in the NICU/SCBU																	
Gentamycin	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
3G Cephalosporins	✓	✓	✗	✗	✗	✗	✓	✓	✗	✓	✓	✓	✗	✓	✗	✓	✓
Metronidazole	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Cloxacillin (PO/IV)	✓	✓	✓	✓	✓	✓	✓	✓		✗	✓	✓	✓	✓	✓	✓	✓
Aminophillin	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Adrenaline	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Fruzemide	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
5% Dextrose	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
10%Dextrose	✗	✓	✓	✓	✓	✗	✗	✗	✓	✓	✓	✗	✗	✓	✗	✗	✗
Nebulisation solutions - Ipravent	✗	✓	✗	✓	✓	✗	✗	✓	✓	✗	✗	✗	✓	✗	✗	✗	✓
Hydrocortizone	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Dexamethazone	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Dopamin	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Dobutamin	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Digoxine	✓	✓	✓	✓	✓	✗	✗	✓	✓	✓	✓	✗	✓	✗	✗	✗	✓
Phenobarbitone (PO)	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✗	✓	✓	✓
Phenobarbitone (IV)	✓	✓	✓	✓	✓	✓	✗	✓	✗	✓	✓	✓	✓	✓	✓	✗	✓
MgSO4	✓	✓	✗	✓	✓	✓	✗	✓	✓	✗	✓	✗	✓	✗	✓	✗	✓
Sildenafil	✗	✗	✗	✗	✓	✓	✗	✗	✗	✗	✗	✗	✗	✗	✓	✗	✗
Glucagon	✓	✗	✗	✓	✓	✗	✗	✓	✓	✗	✗	✗	✗	✗	✗	✗	✓
NaHCO3	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
KCl	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Having a surfactant vial in stock	✓	✗	✗	✗	✗	✓	✓	✓	✓	✓	✓	✓	✗	✗	✓	✓	✓
Ability to local purchase surfactant when needed	✓	✓	✗	✗	✗	✗		✓	✓	✓	✗	✗	✗	✗	✗		✓

Table 10.12: Facilities for infection control by institution and district

	COLOMBO									GAMPAHA					KALUTARA		
	TH CSHW	LRH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTIWALA	DGH KALUTARA	BH PANADURA	BH HORANA
UFI	L101		L102	L102	L102	L104	L104	L105	L106	B101	B101	B302	B203	B304			
FACILITIES FOR INFECTION PREVENTION																	
Soap	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓
Elbow taps	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Running water	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Hand rub solutions	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Gloves	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓
Isolation facilities	✓	✓	✗	✗	✗	✓	✗	✓	✗	✗	✓	✓	✗	✓	✓	✗	✓
Disposable towels (single use)	✓	✓	✗	✓	✗	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓
Hand washing facility available at the entrance	✗	✓	✓	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗
A hand rub solution bottle per each section	✓	✓	✓	✓	✗	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
The hand washing/ hand rub practice satisfactory	✓	✓	✓	✓		✗	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓

Some of the basic facilities for infection control such as elbow taps, gloves, single use disposable towels, hand rub solution bottle per each section are not available in some of the institutions in the Southern Province. These are extremely cost effective practices for the prevention of infections in the newborn.

This table depict the availability or non-availability of essential equipment in the SCBUs and NICUs in the institutions in the Western Province. However the adequacy of essential equipment according to the case load is not calculated here.

Table 10.13 indicate the standard equipment that should be available in the NICUs and SCBUs.

Table 10.13 Equipment available at NICUs and SCBUs by institution and district

	COLOMBO									GAMPAHA					KALUTARA		
	TH CSHW	LRH	TH DMH	TH DMH	TH DMH	TH KALUBOWILA	TH KALUBOWILA	BH AVISSAWELLA	BH HOMAGAMA	TH RAGAMA	TH RAGAMA	DGH GAMPAHA	DGH NEGOMBO	BH WATHUPITTWALA	DGH KALUTARA	BH PANADURA	BH HORANA
UFI	L101		L102	L102	L102	L104	L104	L105	L106	B101	B101	B302	B203	B304			
EQUIPMENT AND SUPPLIES																	
Multipara monitors	✓	✓	✗	✗	✓	✓	✗	✓	✓	✓	✓	✗	✓	✓	✗	✓	✓
Pulse oxymeters	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓
Open resuscitation table with overhead warmer	✓	✓	✓	✓	✗	✓	✗	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓
Neonatal resuscitation trolley	✓	✓	✓	✓	✗	✓	✗	✗	✓	✗	✓	✓	✓	✓	✓	✓	✓
Syringe pumps	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Infusion pumps	✓	✓	✗	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Nebulizers (Oxygen driven)	✓	✓	✗	✓	✓	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Electronic weighing scale (Digital)	✗	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗
Portable sucker (one jar)	✓	✓	✗	✓	✗	✗	✗	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓
Cold light for IV cannulation	✗	✓	✗	✗	✗	✗	✓	✓	✓	✗	✓	✓	✓	✗	✓	✓	✗
Spot lamp	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓
Computer	✓	✗	✗	✗	✗	✗	✗	✗	✗	✗	✗	✓	✗	✓	✓	✗	✓
Central suction	✗	✓	✗	✗	✗	✗	✓	✗	✗	✓	✓	✓	✗	✗	✗	✗	✗
Umbilical probes	✓	✓	✗	✗	✗	✗	✗	✗	✗	✓	✓	✓	✗	✗	✗	✗	✗
Phototherapy units	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Perspex shields	✓	✓	✗	✗	✗	✗	✗	✗	✓	✓	✓	✗	✗	✗	✗	✗	✗
Neonatal stethoscopes	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓
Steel drums	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
IV infusion stands	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
X ray illuminator	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Ward round trolley	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Tapes	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓
Torch	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓
Ophthalmoscope	✓	✓	✓	✗	✓	✓	✓	✗	✓	✓	✗	✓	✗	✓	✓	✗	✓
Charts (Growth, Bilirubin)	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✗	✗	✗	✗	✗	✗
Lab request forms	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓
Calculator	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✗	✗	✗	✗	✗
O2 cylinders with flow meters & humidifier bottle	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Empty O2 cylinders	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓	✓
Head box	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗
Cots	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Incubators	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗	✓	✓
Ventilators	✓	✓	✗	✗	✗	✓	✗	✓	✓	✓	✓	✓	✓	✗	✓	✓	✗
Thermometer	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Umbilical cannulation set	✓	✓	✗	✗	✗	✗	✗	✓	✗	✓	✓	✓	✗	✗	✓	✗	✗
Exchange transfusion set	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Wall dispenser	✓	✓	✗	✗	✗	✗	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Wall clocks	✓	✓	✓	✓	✓	✓	✗	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

Chapter 11

Recommendations

The recommendations presented in this chapter are based on the findings of the assessment. The draft recommendations were presented to the core group and at a national stake holder meeting for discussion.

The nine Provincial reports provide detailed information on individual institution included in the assessment. The tables and figures presented in the provincial reports highlight the deficiencies identified at institutional and district level. Attending to these deficiencies at institutional level will enable achieving the broader recommendations outlined at national level.

11.1 Signal functions

- a) It is recommended that a dialogue is initiated with relevant stakeholders and a policy decision taken as to the signal functions that should be available at the different types of government health facilities.

The following suggestions are presented for initiating a dialog:

- All 9 signal functions must be provided in all institutions which have specialist services (CEmONC) and the aim should be to provide these services 24*7.
- Identify the signal functions that should be available in institutions where specialist

services are not available and define BEmONC in the Sri Lankan context. In this respect, it is noted that evacuation of retained products is not expected to be carried out in institutions without specialist services. Consensus is needed on performing assisted vaginal delivery and manual removal of placenta in such institutions.

11.2 Availability and accessibility

The findings of the survey suggest that internationally recommended standard of 5 EmONC facilities per 500 000 population of which at least one, should provide CEmONC services while the others provide BEmONC services is not appropriate for Sri Lanka. Further, the assessment revealed that 93.1% of deliveries which occur in the government sector health institutions, take place in CEmONC facilities. Therefore, it is important that a standard applicable to the country be agreed upon, taking in to consideration the current birth rate, resource availability and also the people's aspirations and choices regarding type of institution for child birth.

- a) It is recommended that a dialogue be initiated to develop a national standard for availability of EmONC and it is suggested that a minimum of 3 CEmONC facilities per 500,000 population be considered. However, in developing such a standard it is necessary to consider the fact that some of the institutions in Sri Lanka have multiple units

totalling up to 117 specialist maternity units in the country.

- b) Availability has to be considered together with accessibility. It is therefore recommended that geographic location of institutions be such that the total population would fall within a 30 km radius of a 24*7 CEmONC facility.

A time bound plan for achieving the above must be developed and implemented.

11.3 BEmONC facilities

- a) It must be ensured that the signal functions identified as appropriate in the Sri Lankan context for institutions without specialist services are made available in all institutions where women seek care for delivery.
- b) Although only 6.9% births take place in non specialist hospitals, it is important to give due consideration to the high level of direct obstetric morbidity reported from these institutions and the potential to generate nearly 3600 complications per year (from 24000 deliveries).
- c) Special attention must be directed to districts where percentage of deliveries in such institutions is high such as Nuwara Eliya (24%), Mannar (22%), Mullativu (19%), Anuradhapura (16%), Batticaloa (14%) and Monaragala (11%).
- d) It is recommended that Divisional Hospitals for strengthening of maternity services be prioritized taking into consideration the cost effectiveness, utilisation of maternity services and geographic distribution of other institutions in the area.
- e) In order to strengthen services at Divisional Hospitals, it is recommended that norms

for infrastructure and human resources profiles for BEmONC institutions be defined utilising the findings of this assessment (Staff - MO/Nurses/Midwifery qualified nurses/ Midwives and their training, Essential drugs, Equipment and infrastructure facilities)

- f) The need for continuing provision of BEmONC services in the 151 Divisional Hospitals where not even a single delivery was reported in 2011 must be reconsidered. In doing so, the underlying social and economical factors such as poverty, road network, and educational status of women need to be looked at carefully.

11.4 CEmONC and 24*7 CEmONC facilities

- a) It is recommended that norms and standards in respect of infrastructure and human resources necessary for provision of CEmONC and 24*7 CEmONC services be developed for the different types of institutions and suitable arrangements be made to ensure that these standards are adhered to.
- b) It must be ensured that all supporting facilities for the provision of 24*7 CEmONC services are available in institutions where there are 2 or more specialist obstetricians.

An additional 83 specialist obstetricians would be needed to make all single specialist obstetrician (31) stations to two specialist obstetrician stations and to appoint two specialist obstetricians to upgraded BHs without specialist (26 stations) so that all such institutions are capable of 24*7 CEmONC services.

- c) Therefore, it is recommended that this be carried out in a phased manner. Priority stations for appointing a second specialist obstetrician needs to be identified taking into account the population and area to be covered so as to maximise 24*7 CEmONC service coverage.

Table 11.1 (a) Institutions suggested for upgrading in the first phase

Institution for appointment of a second specialist obstetrician	Upgraded BHs to be made functional by appointing specialist obstetricians
DGH Killinochchi	BH Tissamaharama
DGH Mannar	BH Bibile
DGH Mullativu	BH Mahaoya
BH Dambulla	BH Padaviya
BH Nikeweratiya	BH Galgamuwa
BH Mahiyangana	BH Potuvil
BH Kantale	BH Wellawaya
BH Dickoya	BH Deniyaya
BH Balangoda	BH Welimada
BH Valachchenai	

- d) It is recommended that the institutions listed in table 11.1 (a) be strengthened in the first phase to provide 24*7 CEmONC services. Figure 11.1 (a) shows that if the above

hospitals are upgraded, nearly the total area /population of the island will fall within 39 km. radius of a 24*7 EmONC facility.

Figure 11.1 (a): 24*7 CEmONC coverage after upgrading of institutions listed in table 11.1 (a)





Table 11.1 (b) Institutions suggested for upgrading in the second phase

Institution for appointment of a second specialist obstetrician	Upgraded BHs to be made functional by appointing specialist obstetricians
BH Madirigiriya	BH Kalawana
BH Thambuttegama	BH Kabethigollewa
BH Dehiattakandiya	BH Siyambalanduwa
BH Muttur	BH Mankulam
	BH Cheddikulam
	BH Welikanda

- e). It is recommended that the institutions given in table 11.1 (b) be upgraded in the second phase. Figure 11.1 (b) shows that at the end of phase 2 nearly the total population of the island will fall within a 30 km radius of an institution providing 24*7 CEmONC services.

Figure 11.1 (b): 24*7 CEmONC coverage after upgrading of institutions identified for phase two



- a) Since the provision of 2 specialist obstetricians at each of the above institutions would need 28 additional specialist obstetricians in the first instance, until such numbers are trained the following special arrangements are recommended as interim measures:

- **Clustering of institutions**

It is important to explore the possibility to cluster hospitals in such a way that one or more hospitals provide 24*7 services for the cluster.

A second option is to cluster two hospitals which drain relatively small populations together and appoint three specialist obstetricians. This will need special provisions.

Expert consensus is needed to identify institutions that can be clustered together for such covering up arrangements. A written policy and protocol, in the form of an SOP indicating the exact functional details is essential. Transport and communication facilities necessary between the cluster institutions have to be identified. Pilot testing of the project in one or more districts is recommended.

- **Appointment of Senior Registrars**

Explore the possibility of appointing Senior Registrars (post MD pre Board Certification level) to institutions with a single specialist obstetrician. This needs to be done with the concurrence of SLMC so that the appointed officer is able to perform surgery on his own.

- b) The study identified difficulties in providing care in situations where two specialist obstetricians share one ward. Therefore, it is recommended that establishing two independent units be considered in the situations where the monthly deliveries exceed 300 per month.

11.5 Special critical care facilities

It is recommended that special care for obstetric emergencies be made available at three levels.

- a) It is recommended that every specialist unit have 2-4 high dependency beds for pregnant women who need close monitoring. The number of beds needs to be decided based on the case load. The essential equipment for these needs to be defined and provided. Training of nursing staff for close monitoring of patients at HDU can be arranged within the institution/district or provincial level with an ICU.
- b) In the hospitals with ICU facilities, it is recommended to develop and implement admission criteria for obstetric patients in consultation with obstetricians and the specialist in charge of ICUs.
- c) It is recommended that a suitable (web based) mechanism be developed and implemented to identify the closest available ICU beds at any given time in order to ensure a smooth path way of care for these patients without delay.
- d) Highly specialised multidisciplinary critical care facilities at national level in selected tertiary hospitals.

It is recommended that “highly specialised” maternity care units be established in strategic locations with multidisciplinary support teams for the management of emerging issues such as placenta accrete, liver or renal failure secondary to obstetric emergencies which need sophisticated management modalities. It is suggested that at least one such unit be established initially at national level and to increase the number in a phased manner.

In view of the increasing contribution of cardiovascular disease to obstetric morbidity and mortality and availability of advanced treatment modalities which provides an opportunity to manage

seriously ill cardiac patients, it is recommended that an obstetric unit/s be identified and developed to work in collaboration with National Institute of Cardiology / cardiology units to provide such services.

It is recommended that clearly identified protocols and pathways be developed for emergency transfers.

The organisation of rapid response teams within institutions, to deal with emergencies are recommended. These could be started in the higher level institutions in the first instance.

11.6. Establishment of Norms

- a) The study identified high variability between and within districts in resources, especially human resources. It is recommended that norms to be identified for

- Human Resources
- Drugs
- Equipment
- Infrastructure
- Laboratory facilities

based on the workload, type of institution and level of clinical management implemented at each level. A committee needs to be appointed comprising FHB, MoH and professional colleges to finalise the norms and standards.

- b) Malfunctioning equipment was a main reason given for scarcity of equipment and cancellation of theatre lists. Management of equipment to ensure functionality needs to be focussed on at all levels. This can be monitored at institutional level by developing SOPs, introducing check lists and by delegating the responsibility to relevant staff.
- c) Continuous water supply, electricity supply and communication facilities need to be ensured at all level of facilities.

Human Resources

- a) The study findings highlighted the need for clear definition of roles and responsibilities of different levels of staff as well as issues of accountability. It is recommended that clarity in the roles and responsibilities be established with regard to different aspects of care including the management of specific emergencies by way of developing duty lists, guidelines and flowcharts with these details identified.
- b) Task shifting should be considered where necessary.
- c) It is recommended that these be used to establish a culture of accountability and a mechanism for monitoring within the institutions cross cutting all levels of health care providers.

11.7. Care practices

The data collected on selected practices show that there is much variation in the care practices between institutions.

- a) It is recommended the level of complications that can be managed at each level of institution be defined along with clear instructions for decision making and referrals.
- b) In order to achieve high quality and uniformity in maternity care practices it is recommended that the MoH, SLCOG and other relevant professional colleges initiate discussions and reach a consensus on the clinical standards and thresholds for selected care practices.
- c) It is recommended that steps to be taken to institutionalize the use of the partograph at all levels of hospitals.

- d) It is recommended that guidelines for the management of labour, uncomplicated delivery and common complications and emergencies be developed and implemented as to ensure streamlined, effective and uniform care practice.

Suggested priority topics to be considered for such guidelines are:

- Induction of labour
 - Episiotomy (decision, perform, suturing)
 - Management of uncomplicated labour
 - Pain relief during labour
 - Active management of third stage of labour
 - Management of direct obstetric emergencies (especially PPH, eclampsia and pre-eclampsia)
 - Indications for caesarean sections especially for primipara
 - Common complications; shoulder dystocia, cord prolapse, IUGR
 - Post abortion care
- e) It is recommended that step by step procedures (algorithms/ clinical pathways / flowcharts) be developed summarising the above mentioned guidelines and their use ensured through circular directives. These should identify levels of clinical management that should be implemented at each type of institution with clear guidance on referral to higher levels.
- f) It is recommended that a collection of guidelines on the management of obstetric emergencies in non-specialist hospitals be developed in the form of a hand book and disseminated supported by a circular issued by the MoH. They need to be action oriented and specifically designed considering the resources available in these institutions and should include criteria and guidance on transferring such patients to higher level institutions.

- g) It is recommended that Magnesium Sulphate be available at all levels of hospitals where obstetric care is provided. This needs to be supported by the development of a guideline on stock management and use of Magnesium Sulphate in specialist and non-specialist hospitals and disseminated to all institutions through a MoH Circular.

- h) It is recommended that a clearly defined policy for pain relief and companionship during labour be developed by the FHB in consultation with the SLCOG and College of Anaesthesiologists for different levels of hospitals. This needs to be accompanied by efforts to create a “culture of providing pain relief” in health institutions through a process of sensitisation and capacity development of the staff.

11.8. Quality assurance process

- a) Improvement of quality of EmONC care is a priority. A strategy document on quality improvement identifying key activities be developed by the FHB in collaboration with the Directorate of Health Care Quality and Safety, SLCOG, other professional colleges and with care seeker representation.
- b) It is recommended that systems for monitoring quality of care such as clinical audits, SWOT analysis of near misses, adverse incident reporting, clinical quality circles and tools such as the maternity dashboard be developed. Such systems should be planned and implemented jointly by the FHB and the SLCOG in collaboration with the Directorate of Health Care Quality and Safety.
- c) It is recommended that FHB in collaboration with Directorate of Health Care Quality and Safety and professional colleges reach consensus on clinical quality of care indicators and these be included in the National Quality Assurance Programme. A

set of key indicators also be developed to facilitate the in-house monitoring by heads of institutions (through heads of units meetings) to address the gaps in quality.

- d) Explore the possibility of developing a system of supportive supervision of the hospitals at each level to ensure that the quality assurance measures are being implemented.

11.9. Caesarean Sections

The findings on caesarean sections merit wider discussion at a professional forum.

- a) In order to generate valid evidence for such an exercise, it is recommended that a national in-depth study be conducted on the trends, both health and non-health determinants, care provision and a cost assessment of caesarean deliveries in the country including those in the private sector. The study should also aim to address the reasons for the high variability in CS rates between institutions. It is also recommended to conduct a secondary analysis of the WHO Multi Country Survey on Maternal and Newborn Health.
- b) It is recommended that the findings of such studies be converted in to progressive action by the united efforts and commitment of the SLCOG, FHB and MoH through Technical Advisory Committee on Maternal Health and Family Planning and National Committee on Family Health.

11.10 Maternal Mortality and Morbidity

- a) Obstetric haemorrhage still accounts for a large proportion of the maternal deaths. It is recommended that every effort be made to ensure high quality blood transfusion services including 24*7 cover by a medical officer from the NBTS. As an interim measure an adequately trained and competent, designated MO other than ward HO/MOs be made available.

- b) It is recommended that a culture of a PPH drill in all obstetric units be developed and encouraged through routine skills development activities. Measurements need to be taken to ensure that a “PPH management trolley” be available with necessary drugs, equipment and supplies for immediate response.
- c) It is recommended that the country change to a confidential inquiry into maternal deaths. This should be a well-planned transition from current practice with adequate consultations with relevant stakeholders and learning from the experiences of other countries where successful reviews are conducted.
- d) It is recommended that inquiry in to “near misses” be instituted early. In the current scenario of decreasing mortality this is essential for the identification of points at which service provision may be improved. This is also likely to be less sensitive than inquiry into mortality since the focus is on deaths averted. Data collection on this aspect may be piloted in selected sentinel institutions / districts and scaled up to national level in a phased manner.
- e) The information from the pilot phase may be used to validate/refine the criteria / definitions used in identifying “Severe Maternal Morbidity”.
- f) Once the “near miss” inquiry is established nationally, it is recommended that severe maternal morbidity indicators such as Severe Maternal Outcome Ratio (SMOR), Maternal Severity Index (MSI), prevalence of women with life threatening conditions (WLTC), Maternal Near miss Mortality Ratio, (ratio between maternal near-miss cases and maternal deaths) , be accepted as quality of care indicators at national and subnational level and be reported routinely in addition to the MMR.

11.11. Supportive services for EmONC

- a) It is recommended that public health, health education and infection control units be established in all hospitals with specialist services, defining specific roles and responsibilities.
- b) It is recommended that dedicated obstetric operating theatres based on the case load be established in selected institutions and explore the possibility of having a designated theatre table for obstetric emergencies in other institutions.
- c) It is recommended that 24*7 laboratory services be available in all specialist hospitals. In the interim period it is necessary to consider making available selected investigations necessary for management of emergencies on a 24*7 basis.

11.12. Training

- a) It is recommended that in service-training needs for the different categories of staff be revisited and the training be competency based. Training plans and modules should be agreed upon and prepared nationally, so that irrespective of the training agency, there would be uniformity of messages. The plans and modules need to be revised every 5 years.
- b) The survey noted that use of the partograph and recording of events, both were poor especially in non-specialist institutions. A module that would impart knowledge and skills necessary to monitor labour using the partograph and a plan to improve its utilisation is important. The WHO self-learning module may be used for this purpose. In addition to the obstetric topics, in-service training modules should include the following; family planning, effective communication, basics of counselling and befriending.

- c) In conducting training, the staff should be trained as teams comprising doctors, nurses and midwives and should include emergency drills.
- d) Skills laboratories can be established in the selected hospitals /regional training centres/ districts in order to ensure competency based training. MOMCH or MO/Training at RTC can coordinate the training and mobilize local resources.

The training teams for districts can be trained at national level using the modules developed.

- e) It is recommended that a cluster system be established within a district/province for technical guidance, training, and to ensure clinical management based on nationally identified protocols and guidelines. It is recommended that specialist obstetricians working in the district should lead this process.
- f) The FHB with the professional colleges and quality secretariat need to establish a system for monitoring the quality of training at subnational level.
- g) It is recommended that each hospital / district maintain an easily accessible in-service training profile of its staff and that the district have a fixed pre planned training schedule and a budget enabling institutions to maximise the training opportunities.

11.13. Institution based family planning services

- a) Mortality and morbidity due to septic abortion remains high. Thus, minimising unwanted/ unplanned pregnancies is a priority. Delay in seeking care for septic abortion was noted, as such it is important to decriminalise the abortion seeker and improve the quality of post abortion care especially in the area of interactions with care providers.

- b) It is recommended that the institutional based family planning services be re-organised nationally. The services and the frequency with which they have to be provided by the different types of institutions be identified clearly. Ideally the hospitals with CEmONC facilities should conduct family planning clinics on 7 days a week.
- c) The roles and responsibilities of Obstetric and Gynecology ward staff in providing family planning services have to be clearly identified and circularised.
- d) Dedicated theatre time for the provision of permanent methods have to be identified at institution level.
- e) Stock outs of all family planning commodities were noted, as such it is recommended that stock control procedures be re-organised and re-instituted.
- f) It is recommended that the flow of information from the institution based FP clinics to the national health information system has to be strengthened.
- g) It is recommended that mechanisms for monitoring the implementation of these activities at district level have to be strengthened.
- h) Regular in-service training and updating knowledge on family planning should be made available for institutional staff.
- i) The post- partum IUD insertion to be piloted in selected institutions

This should be arrived through a consultative process that includes all relevant stakeholders.

The following levels are suggested for discussion towards a national consensus:

(1) Basic Neonatal Care (Essential Newborn Care)

It is recommended that basic neonatal care (Essential Newborn Care) is made available in all institutions that provide maternity care.

They should be able to provide the following services;

- Immediate care of the newborn at the time of delivery,
- Thermal care,
- Prevention and control of infections,
- Basic neonatal resuscitation,
- Support for breastfeeding,
- Examination of the newborn,
- Stabilization and provide care for newborns needing special care prior to transfer to a facility that can provide the appropriate level of neonatal care.

(2) Higher levels of care

It is recommended that special care and intensive care for the newborns be categorized under four levels.

Levels I and II would provide special care graded in a hierarchical fashion.

Levels III and III + would provide neonatal intensive care.

11.14 Neonatal care services

11.14.1 Availability and accessibility

- a) It is recommended that a policy decision be made regarding the different levels of neonatal care in the country as well as the services to be provided at each level.

A hierarchical structure is recommended taking in to consideration the fact that provision of highly specialised neonatal care is effective only when an adequate number of cases are managed at a given time. So that the care providers are able to develop their expertise in management. Furthermore, equipment and infrastructure needed for special care

is expensive and the provision of services need highly skilled personnel.

The following levels of care are recommended:

Level I neonatal care (special care)

It is recommended that all base hospitals be provided with Level I neonatal care.

Level II neonatal care (special care)

It is recommended that each district general hospital be provided with facilities for this level of care.

Level III neonatal intensive /critical care

This level would provide neonatal intensive care and would be limited to THs and PGHs. It is recommended that at least one facility is available per province.

Level III + neonatal intensive care

This would be limited to a few (3-4 for the country) special teaching hospitals where highly specialized services would be available such as neonatal cardiac and surgical facilities and the management of babies born with an extreme low birth weight (<1500 gms). or born at around 22-24 weeks of gestation.

It is recommended that the Lady Ridgeway Hospital, Teaching Hospital Karapitiya and Sirimavo Bandaranayake Children's Hospital and the TH Jaffna/Vavauniya/Anuradhapura be considered for a level III + institutions.

- b) It is recommended that the establishment of neonatal intensive care be done in a phased manner with the establishment of one or two Level III + facilities in the first phase.
- c) In deciding on the phasing of development of other levels of care it is recommended that institutions be selected based on the following criteria:

Population (births) as well as the area to be served

Terrain to be covered

Proximity to other facilities/ other level facilities

Time taken for transfer/retrieval between institutions

High neonatal mortality

Social determinants such as poverty, educational level etc

11.14.2. Establishment of Norms

- a) The study identified high variability between and within districts in resources.

It is recommended that norms to be identified for

- Human Resources
- Drugs
- Equipment
- Infrastructure

based on the level of care, workload and the type of institution.

A committee needs to be appointed comprising FHB, MoH and professional colleges to finalise the norms for neonatal care.

- b) Management of equipment to ensure functionality needs to be focused at all levels. This can be monitored at institutional level by developing SOPs, introducing check lists and by delegating the responsibility to relevant staff.
- c) Continuous water supply, electricity supply and communication facilities need to be ensured at all level of facilities.

11.14.3 Human Resource

- a) Study findings clearly demonstrate that it is necessary to identify norms on human resource for the different levels of neonatal care (Level I, II, III or III+) so as to ensure 24*7 services.
- b) It is recommended that Neonatologists are appointed to all the Level III and III+ institutions. It is recommended that human resources necessary to provide 24*7 services be available.
- c) Study findings indicate that some of the institutions cannot provide 24*7 neonatal intensive care due to lack of adequate number of medical officers. Designated medical officers for neonatology has to be appointed to all the level III and III+ institutions and to level II institutions based on workload. Norms for medical officers should be developed taking into consideration the work load, and the need to provide 24*7 service provision by this category of MO.

11.14.3 (b) Training

- a) It is recommended that the training requirements for different categories of personnel be identified according to the categorisation of care levels given above.
- b) Norms/standards for training of the staff involved in neonatal care be established.
- c) District training pools need to be developed and budgetary allocations provided for training.
- d) Training centres to provide such training has to be identified and training should be organised according to a schedule. Already established training programmes for neonatal care such as the Essential Newborn Care Course, Neonatal Advanced Life Support Course, 40 hours Breastfeeding

Counselling Course, Baby Friendly Hospital Initiative Course and the Baby Friendly Hospital Initiative Course for the Administrators have to be regularly conducted according to a set plan so as to cover all the staff caring for newborns.

- e) It is recommended that the in service training module on Care of the Sick Newborn developed under the SDF/ MCH project has be adapted and used for training.

11.14.4 Clinical standards and quality of care

The data collected on selected practices show that there is much variation in the care practices between institutions.

- a) It is recommended the level of management of complications that can be carried out at each level of hospital be defined along with clear instructions for decision making and referrals.
- b) Standards for newborn care have to be introduced. Systems for clinical auditing and quality assurance have to be developed and introduced to all levels of institutions.
- c) It is recommended that guidelines for care of the sick neonate have to be prepared and introduced to the institutions and regular use ensured to achieve quality assurance in clinical care.

Suggested priority topics to be considered for such guidelines are:

Care of normal newborn at birth and beyond
Hypothermia and thermal control
Breastfeeding
Management of LBW babies
Management of hypoglycaemia
Respiratory distress in newborn
Neonatal jaundice

Neonatal sepsis

Emergency triage assessment and treatment

Neonatal transport

- d) It is recommended that step by step procedures (algorithms/ clinical pathways / flowcharts) be developed summarising the above mentioned guidelines and their use ensured through circular directives.
- e) It is recommended that the Neonatal Death Investigation be streamlined including the Perinatal Death Audit. Mechanisms for monitoring follow up actions have to be formalised. .

11.14.5 Referral system

- a) It is recommended that protocols and guidelines be developed for the following:
- Referral
 - Accepting referrals
 - Referral pathways and
 - Retrieval of sick newborns
- b) It is recommended that in-utero transfers be sent to a Level III centre as much as possible. After delivery if no complications are identified they may be transferred back to the hospital of origin.
- c) SOPs should be developed on transfer of newborns.

11.14.6 Neonatal transport system

- a) With the development of the proposed levels of care, it would be essential to establish a well-structured neonatal transport system. It is recommended that a dedicated neonatal transport system be established with trained retrieval teams at designated centres. A centrally located bed management system is recommended linking all the neonatal units in the country.

11.14.7. Mother Baby Centres / Lactation Management Centres

- a) It is recommended that Mother Baby Centers and Lactation Management Centers be established as per current MoH recommendation; i.e. they should be available in all the hospitals providing specialist care: from Base Hospitals upwards i.e. in all institutions providing special care and intensive neonatal care.

11.15 Others

- a) The survey findings suggest that pharmacy management could be improved to maximise the effective utilisation of drugs.
- b) Poor documentation practices were encountered at all levels. Entering of diagnosis in BHTs, registers, records and operation notes in the ward and theatres need much improvement. Reporting of still births were poor and this needs immediate attention.
- c) It is recommended to introduce ICD 10th classification into hospital Morbidity and Mortality recording and return system thus allowing international comparisons.
- d) In order to overcome the undue delays in compilation of hospital statistics, it is recommended to introduce electronic IMMR on a phase basis.
- e) Steps should be taken to introduce an obstetric MIS. It may be piloted in one or two districts with a view to cover the whole country later.
- f) It is recommended that information on neonatal deaths within the first 24 hours of life be included in the routine MIS.

REFERENCES

Department of Census and Statistics. 2012. Census of population and housing 2012 [Online]. Department of Census and Statistics. [Accessed January 2013].

Family Health Bureau of Ministry of Healthcare and Nutrition, Perinatal Society of Sri Lanka & Central Engineering Consultancy Bureau 2007. Building and other guidelines for Neonatal Intensive Care Units, Special Care Baby Units, Mother Baby Centres.

Registrar General Department. 2012. Birth registration [Online]. Registrar General Department. [Accessed January 2013].

United Nations 1997. Guidelines for Monitoring the Availability and Use of Obstetric Services.

World Health Organization 2009. Monitoring Emergency Obstetric Care: A handbook. Geneva.

Annexure 1: Calculation of EmONC indicators

Indicator	Description	How it is calculated	
1	Availability of emergency obstetric care: basic and comprehensive care facilities	Numerator	No of facilities in area providing basic or comprehensive EmONC
		Denominator	Population of area ¹ divided by 500 000
2	Geographical distribution of emergency obstetric care facilities	Numerator	No of facilities in area providing basic or comprehensive EmONC
		Denominator	Geographic extent of area in km ²
3	Proportion of all births in emergency obstetric care facilities	Numerator	No. of women giving birth in EmONC facilities in a given period
		Denominator	Total no. of births reported in area for the same period
4	Meeting the need for emergency obstetric care: proportion of women with major direct obstetric complications who are treated in such facilities	Numerator	No. of women with major direct obstetric complications treated at EmONC facilities in a specified period
		Denominator	Expected no. of women with severe direct obstetric complications in area for the same period (15% of reported births in the same area and period)
5	Caesarean section as proportion of all births	Numerator	No. of caesarean sections in EmONC facilities in specified period
		Denominator	Total no. of births reported in area for the same period
6	Direct obstetric case fatality rate	Numerator	No. of maternal deaths due to direct obstetric causes in EmONC facilities in specified period
		Denominator	Estimated no. of women treated for direct obstetric complications in EmONC facilities in same period
7	Still-birth death rate	Numerator	No. of stillbirths in EmONC facilities in specified period
		Denominator	No. of women giving birth in EmONC facilities in same period
8	Proportion of maternal deaths due to indirect causes in emergency obstetric care facilities	Numerator	No. of maternal deaths due to indirect causes in EmONC facilities in specified period
		Denominator	All maternal deaths (direct and indirect) in EmONC facilities in same period

